

Myles Allan, Constraints for Probabilistic Forecasts

Myles Allan,

Climate Dynamics Group, Atmospheric, Oceanic and Planetary Physics,
Department of Physics, University of Oxford, United Kingdom

This talk will review some basic issues in the use of observational constraints for the generation of probabilistic forecasts on climate (multi-decadal) timescales. Interesting issues arise from the fact that we are making a prediction of an unprecedented, one-off event, leading to questions about what constitutes verification of a probabilistic forecast.

I will introduce the notion of STAID (Stable Inference from Data) forecasts as a reasonable aspiration, and illustrate how forecasts of some very large scale climate variables may be on their way to being STAID already, but formidable challenges lie ahead before we can bring this methodology down to a spatial scale small enough for most practical impact assessments.

I Ansorge, Topographical steering of the Antarctic Circumpolar Current in the south-west Indian Ocean

I Ansorge

Oceanography, University of Cape Town, South Africa

No Abstract Information Available

A Barrable, GCM Sensitivity for Climate Change Studies at a Regional Scale in the Swartland, Western Cape, South Africa.

A Barrable 

Climate Systems Analysis Group, Environmental and Geographical Science Department,
University of Cape Town, South Africa

and

Prof Bruce Hewitson,

Climate change is a critical issue for the farmers in southern Africa, especially in the context of inter- and intra-annual climate variability. In particular an examination of future climate change needs to be assessed in view of the sector of vulnerability. For the agricultural sector, this critically includes a regional study of climate change to enable comparative results towards impacts on soil erosion and land degradation. Within this framework, evaluating the effectiveness of the climate models to be utilised needs to be addressed before the impact of the projected future climate can be properly assessed for the region. This study concerns itself with the southwestern Cape region of the Swartland, where wheat farmers are for the most part reliant on the timing and quantity of rainfall to produce viable yields.

An assessment is made of the performance of six global climate models for the southwestern Cape: the CCCMA, CSIRO-Mk2, ECHAM4 / OPCY3, HadCM3, NCARCSM and NCARPCM. By examining the variables of precipitation, temperature and wind speed, and comparing these to NCEP reanalysis data, one may identify the degree to which the model simulations reflect the real climate system processes. The model data used comprises 30 years of simulated climate representing present day conditions based on the A2 SRES scenario of greenhouse gas concentrations. In this way the sensitivity of the model output to the observed climate is gauged, thus establishing which models are applicable to the study region.

Judit Bartholy ,Comparisional Analysis Of Tendencies Of Extreme Climate Indices On The European And The Carpathian Basin Scales

Judit Bartholy 

Department of Meteorology, Eotvos Lorand University, Hungary

and

Rita Pongracz

Since human and natural systems may be especially affected by changes of extreme climate events, the main objective of our research is to detect the possible changes of intensity and frequency of these extreme events. Several climate extreme indices have been analysed and compared for Europe and the Carpathian Basin for the twentieth century based on the guidelines suggested by the joint WMO-CCI/CLIVAR Working Group on climate change detection. These climate extreme indices include the numbers of severe cold days, winter days, frost days, cold days, warm days, summer days, hot days, extremely hot days, cold nights, warm nights, the intra-annual extreme temperature range, the heat wave duration, the growing season length, the number of wet days (using several threshold values defining extremes), the maximum number of consecutive dry days, the highest 1-day precipitation amount, the greatest 5-day rainfall total, the annual fraction due to extreme precipitation events, etc. Therefore, daily maximum, minimum and mean temperature observations and daily precipitation amounts have been used in the present statistical analysis. Our results suggest that similarly to the global and continental trends, regional temperature of the Carpathian Basin has become warmer during the second half of the twentieth century. Furthermore, regional intensity and frequency of extreme precipitation has increased, while the total precipitation has decreased in the Carpathian Basin and the mean climate has become drier.

Swadhin Behera, Modes Of Variability In The Indian Ocean: Statistical Perceptions

Swadhin Behera 

Frontier Research System For Global Change, Japan

and

Prof. Toshio Yamagata

Reorganization of independent physical modes is an important aspect of climate studies. Statistical methods such as EOF technique have provided scope to grasp such meaningful information from massive climate data. In the tropical Pacific, the El Niño/Southern Oscillation (ENSO) signal is captured as the most dominant mode in eigen techniques. The statistical analyses here capture the coupled phenomenon which is shown to influence the global climate variability. In the tropical Indian Ocean, a basin-wide uniform polarity pattern in the sea surface temperature (SST) anomaly is present as the most dominant interannual mode. This basin-wide uniform mode shows a high correlation with the Pacific variability: The peak correlation coefficient of 0.8 is found when the Niño-3 index leads the basin-wide uniform mode by 4 months. Because of the dominant Pacific influence, the tropical Indian Ocean has been mostly considered to be passive in the climate studies. However, recent progress in the Indian Ocean research has revealed a coupled mode widely known as the Indian Ocean Dipole (IOD). The east-west see-saw in the SST anomalies during an IOD event characterizes the phenomenon. Interestingly, this SST signal is captured as the second dominant mode in the EOF analysis. The less frequent occurrence compared to the basin-wide uniform mode provides IOD the second most dominant seat in the EOF analysis. Nevertheless, the IOD has wide societal impact during the year of occurrences. Although simple statistical methods like EOF technique provide important clues of climate variability, sophisticated statistical analyses sometimes are misleading. For example, the IOD which is captured in the EOF analysis and simple composite technique is missed by the VARIMAX analysis. It is shown that the presence of the basin-wide uniform mode misleads such sophisticated analyses: it is necessary to remove the dominant mode to capture the IOD in these methods. It is important that statistical inferences must be supported by our physical understanding for any meaningful insight of the climate system.

Roman Bekryaev, Estimation Of Climatic Trends Significance

Roman Bekryaev

Arctic And Antarctic Research Institute, St.-Petersburg State University, Russia

Estimation of the climatic trends significance is still not completely resolved problem. The completeness follows from the serial correlations of the time series. The serial correlation in turn is connected to the dominated low-frequency climate variability resulting mostly from the chaotic dynamics of climate systems. Using the traditional statistical approaches (that don't take into account that serial correlation) can lead to completely wrong results and conclusions. In the proposed research we explore the climatic trends variance as a variance of generalized stochastic integral (stochastic integral from the multiplication of random function and weight function). Using the convolution of the weight functions we got the expression applicable for the calculation of the variance of generalized stochastic integrals. We have considered as a null hypothesis the suggestion that the "real" climatic trend is absent and the trend estimated from the data set is connected to the limited sample of the stationary random process. As a result the sampling trend has the mathematical expectation equal to zero but the value of variance can be essential. Using the suggestion about the Gaussian random distribution of the air surface temperature we have constructed the density distribution function of the coefficients of linear and parabolic regression. We have analysed some climatic air surface temperature time series. For the Northern Hemisphere air surface temperature from 1901 to 2001 we have found approximately 95-98% trend significance level. In the same time the trend significance level estimated on the base of routine procedure is more then 99.9%. Our results still support the existing global warming concept.

Rasmus Benestad, Record-Value Statistics And Test Of Independent And Equal Distribution (Iid)

Rasmus Benestad 

The Norwegian Meteorological Institute, Norway

It is common to use probability distribution functions (p.d.f.s) in climate research to describe various climatic parameters. The use of p.d.f.s assume the condition that the values are independent and identically distributed (iid) random values. An ongoing climate change implies a non-stationary situation where some of the climatic parameters do not satisfy the iid condition. A test for whether a climatic element satisfies the iid condition is suggested, with an emphasis on the extreme tails of the distribution. A simple expression can be derived for how many times a new record is set in a series of iid random values, and the validity of the null-hypothesis of iid can be easily tested for any observation. A few examples of the application of record-value statistics to testing of stationarity will be given.

AF Beraki, CLIMATE CHANGE SIMULATIONS OVER ERITREA USING THE PRECIS REGIONAL CLIMATE MODEL SYSTEM

AF Beraki, 

University of Pretoria, South Africa

and

CJdeW Rautenbach
FA Engelbrecht

Eritrea is a relatively young country that gained independence in 1991. It is also regarded as one of the developing African nations that appear to be most vulnerable to climate change and variability because of widespread poverty - presumably as a result of poor management practices and a lack of expertise. Previous studies have indicated that countries in the Sahelian latitudes are noticeably affected by climate variability (especially droughts) related to the El Niño Southern Oscillation (ENSO) phenomena. Eritrea is located in the eastern part of the Sahelian latitudes, and is in general regarded as a semi-arid country. Future changes in the climate of this region might influence temperature and moisture prone activities, especially those related to agriculture. The assessment of past regional climate trends and the development of scenarios for possible future climate changes are regarded as important. To address these issues the PRECIS Regional Climate Model (RCM) System from the UK was implemented over the region spanning 5°N - 51°N and 2.5°E - 28°E. The model was firstly forced at its lateral boundaries by a 30-year present-day (1961-1990) integration of the “atmosphere only” HadCM3H global General Circulation Model (GCM). Secondly, the model was forced by “fully coupled” HadCM3 (for sea surface temperatures and sea ice) and “atmosphere alone” HadAM3H GCM derived boundary conditions for the A2 and B2 SRES scenarios in an attempt to generate a 20-year future climate change integration (2070-2090). The experiment allows for verification of present-day spatial and temporal performance of PRECIS over Eritrea, with specific emphasis on temperature and moisture. Possible climate change scenarios are also investigated.

Acknowledgement: Met Office, Hadley Centre for Climate Prediction and Research, UK

Carel N. Bezuidenhout, Assessing The Value Of The Three Month Rainfall Outlook (SAWS) Within An Operational System To Forecast Sugarcane Production In South Africa

Carel N. Bezuidenhout 
University Of KwaZulu-Natal, South Africa

On average the South African sugar industry produces two million tons of sugar per annum, making it one of the ten largest cane sugar producers in the world. Climate variability in the eastern parts of South Africa has led to a 25% coefficient of variance in sugar production over the past ten years. Timely and accurate yield forecasts prior to and during the milling season holds the potential to improve various activities, such as milling operations, international trade and agronomic optimisation, within the industry. A yield forecasting system using a crop model was developed. The system uses climate data to date and substitutes the future with up to ten different analogue years from the history. Analogue years are selected using the SAWS three month rainfall outlook. The aim of this study was to assess the accuracy of yield forecasts when using the SAWS outlook as opposed to a neutral outlook over the period 1998 to 2002. Errors in yield forecasts (on mill and industry scales) issued between January and May were reduced by between 10% and 20% when using the SAWS rainfall outlook as opposed to a neutral outlook. However, yield forecasts based on the SAWS outlook that were issued in September were often inferior. Spatially it seems as if some areas benefit more from the SAWS outlook than others. Nevertheless, there is significant evidence that in general the SAWS outlook enhanced yield forecasts in the sugar industry. The results also suggest that the technique of selecting ten historic analogue seasons might need some revision.

T.A Bigala, ACID RAIN DEPOSITION OVER SOUTH AFRICA

T.A Bigala 

Climatology Research Group, University of the Witwatersrand, South Africa

and

S.J Piketh

M Barenbrug

MT Freiman

Aerosols affect climate by absorbing or reflecting incoming solar radiation, and by affecting cloud microphysics, cloud albedo and precipitation. The Highveld regions of Mpumalanga and Gauteng are the major industrial areas in South Africa. These areas play a significant role in the production and transport of aerosols. The urban and industrial areas in these provinces are important source emitters of sulphur, carbon, and their corresponding oxides, nitrous oxide and nitrogen dioxide. Measurements of aerosol optical thickness from the ground and aerosol size characteristics over the Highveld region were collected using hand-held haze meters and a CIMEL sun photometer. The optical properties of the industrial and urban aerosols over the Highveld region of South Africa were analysed between the winter months of May-June 2002 and 2003 and the summer months of October-November 2002. The aerosol optical thickness observed over Pretoria, Bethal and Marble Hall in winter 2002 was high whose values ranged from 0.05 to 0.7 as compared to Secunda (0.05 to 0.2). In summer 2002, Marble Hall showed aerosol loading of 0.09 to 0.6 above Pretoria, Secunda and Bethal (0.12 to 0.59). In winter 2003 aerosol loadings showed fairly similar results for Sebokeng, Rustenburg and Standerton, with values ranging from 0.05 to 0.5. The Angström exponent parameter had a wide range of 0.1 to 2.5. This indicates a wide range in particle sizes over the Highveld region. Data from the sites over the Highveld has revealed values whose calculations are derived because of the influences of aeolian dust, coarse mode industrial and fine aerosols.

Mr. Albert Boehm, Proper Verification Scores For Areal Probability Forecasts

Mr. Albert Boehm

Boehm Music Studio, USA

Murphy (1973) defined a proper score as one that rewards a forecaster who honestly communicates their degree of belief. Hedging is said to occur when the stated forecast does not correspond to this degree of belief. Murphy was specifically addressing subjective probability forecasts, but the principle holds for objective automated forecasts as well as categorical forecasts. The concept of a proper score can be extended in three ways. First, there is an important transformation between a degree of belief and a probability. For example, the belief that the 99% heaviest rain will occur generally does not correspond to a statement that the probability of rain over 5 cm is 99%. This transformation corresponds to the probability verification term calibration. The degree of belief ranking corresponds to sharpness. A score can be proper with respect to calibration or sharpness, both, or neither. In one case, a forecaster could believe a 0% probability but call it 75% to improve the calibration score of the 70 to 80% bin which had 3 hits and no misses. Next, Murphy used the term hedging to specify an undesirable characteristic. Aside from the misstatement of degree of belief, hedging implies waffling, a vagueness or evasiveness. However, hedging also means to optimize rewards by allocating resources for diverse alternatives - a very desirable characteristic in decision making under uncertainty. Instead improper influence can be used for a score that encourages misstatement. Thirdly, Murphy defined a proper score with respect to expected value. However, other attributes of a score can have improper influence. For example, a large sampling distribution can include many poor scores even with a good expected value. These three effects are exacerbated in areal probability forecasts. Locations in an area can have different probabilities of, say, 5 cm of rain even though the forecaster expects heavy rain at the 99% level over the entire area. A score can be proper for one type of coverage but improper for another. Areas often are highly correlated which inflates the spread of the sampling distribution. Ref: Murphy, A., 1973: Hedging and Skill Scores for Probability Forecasts, J Appl. Meteor., 12, p215-222.

Julian Charles Brimelow, On The Application Of Teleconnection Data To Predict The Severity Of The Alberta Hail Season

Julian Charles Brimelow 

Department Of Earth And Atmospheric Sciences, University Of Alberta, Canada

Alberta is prone to frequent hailstorms during the summer, and the resulting damage amounts to millions of dollars in insured costs annually. It would therefore be advantageous to understand what processes modulate the intensity of the hail season and to then employ this knowledge to predict the potential severity of the hail season in advance. To this end, we analyzed almost 30 years of hail data (from a dense hail observation network) together with composite upper-air maps to identify those large-scale atmospheric characteristics that best differentiated between active and quiescent hail seasons. This analysis revealed that there were indeed marked and consistent differences in the summertime 500 hPa circulation regime over North America between active and quiescent hail seasons. These differences were also reflected in other variables such as the winter snow pack. It has been demonstrated in the scientific literature that the large-scale circulation over North America is modulated by teleconnection patterns, such as the Pacific North American oscillation, with lag times of several months. Therefore, it stands to reason that knowledge of the phase and intensity of certain winter teleconnection indices could be utilized to predict the nature of the summertime large-scale circulation pattern over North America and thereby infer whether the expected circulation patterns will or will not be favourable for severe convection. Correlation map analyses indicated that summertime circulation patterns over western North America are modulated, at least in part, by the Multivariate ENSO index (MEI), Pacific decadal oscillation (PDO) and East Pacific oscillation (EPO) with lead times of up to six months. The teleconnection data were then used to identify thresholds of the MEI, PDO and EPO indices at specific lead times that best differentiated between active and quiescent hail seasons. Our research suggests that active hail seasons typically followed La Nina winters that occurred concurrently with a negative phase of the PDO and EPO. By contrast, quiescent hail seasons typically followed El Nino winters that occurred concurrently with a positive phase of the PDO and EPO. This set of guidelines can be applied to teleconnection indices in the months leading up to the summer to determine the likelihood of an active or quiescent hail season. Farmers could potentially use this information to decide how much crop insurance to buy.

Paula Brown, Temperature Extremes In Southern New Zealand

Paula Brown 

University Of Otago, New Zealand

The economy of southern New Zealand is heavily resource dependent. Many of these resources are strongly influenced by climate variability, whether it be pastoral agriculture, horticulture, viticulture, forestry, winter and summer tourism or hydro-electricity generation. Southern New Zealand has some long temperature records of over 100 years, a large amount of climatic variability and is located in a region of the world where little climate information exists. Daily minimum and maximum temperatures display a lot of variability over decades of measurements. When focusing on extremes it is often difficult to distinguish errors from genuine outliers. Most homogenisation methods utilise monthly means and statistical tests to identify inhomogeneities from reference series. However, the stations with long records in New Zealand have few neighbour stations to aid identification of inhomogeneities, particularly for the period 1852-1970. Periods of parallel temperature observations between station relocation allow compositing of temperatures using probability distribution matching methods of Trewin and Trevitt (1996). The CUMSUM method of Rhodes and Salinger (1993) is also applied to identify inhomogeneities. A preliminary analysis of a homogenised temperature series will be presented.

John Caesar, Creation Of Long-Running, Homogeneous Daily Gridded Global Temperature Datasets

John Caesar 

Met Office, Hadley Centre For Climate Prediction and Research, United Kingdom

and

Lisa Alexander

Simon Brown

Long, homogeneous quality-controlled climate data are particularly important for the study of climate extremes, given that inhomogeneities caused by artificial jumps are likely to be detected as erroneous extreme events. For many reasons, long temporal and large spatial scale daily climate datasets have been available for this kind of study. We describe the creation of new homogeneous gridded daily maximum and minimum temperature datasets based solely on quality controlled station data from freely available data sources including the Global Daily Climatology Network (GDCN), Global Climate Observing System (GCOS) and European Climate Assessment (ECA). The station data were quality controlled to test for outliers and homogeneity and anomalies were calculated so that the data were more spatially coherent, particularly over regions of varying elevation. The anomalies were gridded with an angular-distance weighting technique using correlation decay distances calculated over a number of latitude bands due to the seasonally and spatially varying pattern of inter-station relationships. We have produced daily maximum and minimum temperature datasets covering the period 1950 to 2000 on a 2.5 by 3.75 degree resolution grid, with spatial patterns that are coherent and physically realistic in areas with sufficient data density. The results were also compared with the Hadley Centre atmosphere-only model, HadAM3H. Although the model displays a greater range the gridded observations at the extreme upper and lower ends of the distribution for some areas, the results are generally encouraging.

Hong-Xing Cao, Long Memory Of East-Asian Monsoon And Its Self-Memory Modeling

Hong-Xing Cao

Chinese Academy Of Meteorological Sciences, China

and

Hui-zhong He

Rechard Blender

Neng Shi

The observed series of East-Asian monsoon index for 1873 to 2000 are analyzed with statistics; its trends and climate change in different periods have been revealed too. It is shown that the East-Asian monsoon has been decreased since 1873, and its interdecadal change is significant. We focus on the decadal to centennial changes in East-Asian monsoon, which are connected to a long time memory of the East-Asian monsoon. In order to understand the long time memory of the East-Asian monsoon, the detrended fluctuation analysis (DFA) is applied to the time series $x(i)$ ($i=1, \dots, N$) of East-Asian monsoon. The departure series of $x(i)$ are calculated and the integrated series of $x(i)$ are denoted as $y(k)$. Then $y(k)$ is divided into segments of equal length s and the least-square line fitting the data in each segment, $y_n(k)$, is calculated. The series $y(k)$ is detrended by subtracting the local trend $y_n(k)$, and the root-mean square fluctuations of the detrended series $F(s)$ are computed. $F(s)$ is computed for all 'segment-size' s . If $\log F(s)$ increases linearly with $\log s$, then the slope of the line relating $F(s)$ and s in a log-log scale gives the scaling exponent a_H . According to the random walk theory, the fluctuations $F(s)$ in a given time segment of length s are related to the autocorrelation function $C(s)$. For the long time power law correlations exhibit a power spectra: $C(s) \sim s^{-a_H}$, the fluctuations $F(s)$ increase as a power law: $F(s) \sim s^{a_H}$, $a_H = 1 - a_C$. We have computed the East-Asian monsoon indices with the DFA, The results show that the East-Asian monsoon presents a long time memory; it means the East-Asian monsoon has significant persistence in the yearly series. For winter monsoon, $a_H = 0.55$, $a_C = 0.9$; but for summer monsoon, $a_H = 0.73$, $a_C = 0.54$. It is shown that the difference between long time memories of monsoons in winter and in summer is remarkable. As the weather and the climate over China and East Asia are strongly influenced by the East-Asian monsoon, in particular, in summer, so the long time memory of the summer monsoon hints better predictability for climate change. In order to model the change in East-Asian monsoon, a newly-developed time series technique, namely, the data-based self-memory (DASM) model based on self-memory principle of dynamic system, has been used, with which modeling the East-Asian monsoon indices has been made. The calculating results show the model is capable of simulating and forecasting the climate change in East-Asian monsoon.

S Carter, Identifying General Circulation Model Biases Using Self-Organising Maps

S Carter 

Environmental and Geographical Science Department, University of Cape Town, South Africa

and

Bruce C. Hewitson

General Circulation Models (GCMs) are the major vehicle for assessing global climate change. However, it has been broadly acknowledged that there are many regional biases that have to be considered when using these models. With the use of Self Organising Maps (SOMs) one can characterise the types of circulation that each GCM simulates and compare these archetypal circulation fields to those seen in reality. Two fully coupled ocean-atmosphere GCMs were compared to National Centre for Climate Prediction (NCEP) data, using the later as a proxy for reality. Additionally the relative frequencies for each synoptic state were compared between models and NCEP data, allowing for further biases to be identified.

Lynda Chambers, The Changing Climate Of South-Western Australia

Lynda Chambers 

Bureau Of Meteorology Research Centre, Australia

and

Neville Nicholls

In many regions of Australia there has been the suggestion that the climate has either changed or is in the process of changing. This is particularly evident in the south west of Australia, where rainfall has decreased substantially since the mid-20th century. In 1997 several Australian government agencies formed a partnership with the aim of improving management decisions of climate-dependent industries and environmental management, through improved understanding and definition of interannual and interdecadal climate variability. Six years on, this paper highlights some of the key findings of this research, as well as identifying gaps in our current knowledge.

Richard Chandler, Rainfall Simulation Using Generalised Linear Models

Richard Chandler

Department of Statistical Science, University College London, United Kingdom

and

Chi Yang

We consider the problem of simulating sequences of daily rainfall at a network of sites, in such a way as to reproduce a variety of properties realistically over a range of spatial scales. The properties of interest will vary between applications, but typically will include some measures of 'extreme' rainfall in addition to means, variances, proportions of wet days and autocorrelation structure. Our approach is to fit a Generalized Linear Model (GLM) to raingauge data and, with appropriate incorporation of inter-site dependence structure, to use the GLM to generate simulated sequences. We illustrate the methodology using a dataset from southern England, and show that the GLM is able to reproduce many properties at spatial scales ranging from a single site to 2000km² (the limit of the available data). Having established this, we explore the potential for using the GLM to generate future rainfall scenarios conditioned on climate model outputs.

KE Chiloane, Detection of volatile organic compounds (VOCs) from Cape Town Brown Haze.

KE Chiloane 

University of the Witwatersrand, South Africa

and

S.J Piketh

L.B Otter

Cape Town is among the largest tourist destinations on the continent, resulting in tourism being projected as one of its most important economic growth sectors. But brown haze has a strong degrading effect on visibility which is immediately apparent to the general public and tourists. This "brown haze" is described as a brown-coloured smog found predominantly in the wintertime that occurs mostly from April to September due to strong temperature inversions and windless conditions that can occur during these months. This haze leads to the build-up of pollutants emitted into the atmosphere and is normally most intense in the morning after which it lifts and disperses as the day progresses. The haze is also a cause for concern to residents because of health risks that may accompany the visible air pollution. The project aims to characterise volatile organic compounds (VOCs) associated with anthropogenic adverts that are capable of producing photochemical oxidants by reactions with oxides of nitrogen in the presence of sunlight from the haze. Specific VOCs compounds to be characterised are benzene, toluene, ethyl benzene and xylene (BTEX). Volatile organic compounds (VOCs) samples were collected into canisters for GC analysis, from a South African Weather Service (SAWS) research aircraft, Aerocommander 690A (ZS- JRB) from July 29, 2003 through August 26, 2003. Airborne measurements were also carried out using Fourier Transform Infrared (FTIR) for both brown haze and non-brown haze days. Gas Chromatographic (GC) instrument data will be used to validate the FTIR data. Comparisons of these VOCs with other pollutants (e.g. SO₂, NO₂, ozone, etc) as well their transformations during the day will be investigated at different times of the day. In so doing, the evolution, sources, and constituents of the haze can be determined.

Caio Coelho, Forecast Calibration And Combination: A Simple Bayesian Approach For ENSO

Caio Coelho 

Department of Meteorology, University of Reading, United Kingdom

and

Sergio Pezulli

Magdalena Balmaseda

Francisco Doblas-Reyes

David Stephenson

This study presents a simple approach for combining empirical with raw (i.e. not bias corrected) coupled model ensemble forecasts aiming to provide more skillful interval forecasts of an ENSO index. A Bayesian normal model has been used to combine empirical and raw coupled model December SST Nino-3.4 index forecasts started at the end of the preceding July (5-month lead time). The empirical forecasts were obtained by linear regression between December and the preceding July Nino-3.4 index values over the period 1950-2001. Coupled model ensemble forecasts for the period 1987-99 were provided by ECMWF, as part of the Development of a European Multi-model Ensemble system for seasonal to inTERannual prediction (DEMETER) project. Empirical and raw coupled model ensemble forecasts alone have similar mean absolute error forecast skill score, compared to climatological forecasts, of around 50% over the period 1987-1999. The combined forecast gives an increased skill score of 74% and provides a well-calibrated and reliable estimate of forecast uncertainty. This methodology can also be extended to combine gridded equatorial Pacific SST ensemble forecasts from different coupled models (multimodel approach).

Andrew Colman, Predictions Of Lake Volta Inflow Using Statistical And Dynamical Seasonal Forecast Models

Andrew Colman 

Met Office, Hadley Centre For Climate Prediction And Research, United Kingdom

and

Richard Graham

Results of recent research at the Met Office into the use of seasonal rainfall forecasts to improve long-range prediction of inflow into Lake Volta, Ghana will be described. The work has been carried out in collaboration with the Volta River Authority (VRA), Ghana. By way of introduction the scientific basis for seasonal prediction will be briefly reviewed and Met Office systems for making routine seasonal forecasts to 6-months ahead described (a range of Met Office seasonal forecast products, and an introduction to seasonal forecasting, may be viewed at <http://www.metoffice.com/research/seasonal/>) Hydro-electric power generation from the dam at Akasombo produces approximately 50% of Ghanas electricity and prediction of inflow into the lake, particularly that associated with the June to September rainy season, is a priority in managing subsequent use of the water resource and planning the need for supplementary generation from thermal power stations. The need for reliable inflow prediction has been enhanced by an increase in the frequency of relatively dry years since the dam was commissioned in 1966. VRA currently make routine inflow predictions with regression relationships which use preceding observations of monthly-mean catchment rainfall and river flow as the sole predictors, with no inclusion of predictors based on long-range forecasts of seasonal rainfall. New regressions equations which include input from Met Office seasonal-range rainfall predictions for the Volta catchment have been developed and deliver substantial improvements on the existing methods. Seasonal forecast information used includes input from empirical rainfall prediction models, based on observed large-scale sea surface temperature predictor patterns, and also from real-time integrations of a dynamical coupled ocean-atmosphere model (a version of the Hadley Centre climate model). The new inflow prediction method and its improved performance will be described. Although currently specific to lake Volta, the research has scope for wider application in forecasting water availability in other parts of the globe where seasonal prediction skill is sufficient. In this context, a global assessment of prediction skill for seasonal rainfall will be presented.

Mr Andrew Colman, Predictability Of Extreme Rainfall And Temperature Seasons From Cgcm As Demonstrated By The Demeter Handcasts

Mr Andrew Colman, 

Met Office, Hadley Centre For Climate Prediction And Research, United Kingdom

Extreme seasons for example droughts, hot summers often have a disproportionate impact on society and the economy. Predictions of extreme seasons by the EU funded Demeter project Coupled ocean-atmosphere Global Circulation Model (CGCM) hindcasts have been assessed using a variety of skill measures. Extremes are defined as the outer quint (20%) or outer decile (10%) of the observed PDF. Deterministic forecasts have been assessed using Mean Square Skill Scores (MSSS) and using Gerrity skill scores.

Given that extremes are by definition unlikely and given the uncertain nature of seasonal forecasts, extreme forecasts are an ideal candidate for probability forecasts. Model output has been converted to probabilities using discriminant analysis and other methods and probabilities assessed using the Relative Operating Characteristic (ROC) measure. Skill of forecasts using the Met Office GLObal SEAsonal (GLOSEA) forecast model have been compared with the skill of forecasts produced by combining output from 3 models (Met Office GLOSEA, ECMWF and Meteo France) and of persistence. Extreme rainfall and temperature seasons have been found to be slightly more predictable than ordinary seasons on average with highest skill in the tropics. Examples will be shown where the 3 model forecasts perform better than GLOSEA and vice versa. The dependence of skill on extreme definition, geographical location, skill measure and forecast method will also be discussed.

Claire Davies, Towards High-Resolution Probabilistic Climate Predictions

Claire Davies 

School of Geography and the Environment, University Of Oxford, United Kingdom

and

Mark New
Myles Allen
Dave Stainforth

There are two key limitations to the use of General Circulation Model (GCM) predictions for climate change impact assessment: uncertainties in model physics and the relatively coarse spatial scale of GCM data. We describe a project that will provide a solution to these limitations by making use of data from new efforts in quantifying GCM uncertainties, the ClimatePrediction.net (CPdN) project, in order to generate high-resolution probabilistic climate change predictions. The project will embed empirical downscaling software within the HadCM3 GCM that is running in large-ensemble climate change experiments, distributed through the internet on individual 'volunteer' PCs under the CPdN project. The software will analyse diagnostic fields 'on the fly' to generate and archive time series of predictor variables for the downscaling algorithm. The predictor variables will be returned for archive on CPdN sever nodes, for later input into the downscaling algorithm to generate downscaled predictions for each CPdN run (potentially many tens of thousands of predictions). The downscaled predictions can then be analysed to generate probability distributions for key surface climate variables.

R D Diab, Synthesis of recent tropospheric ozone research over Africa

R D Diab 

University of Kwazulu-Natal, South Africa

A number of large field experiments (for example, SAFARI-92, SAFARI-2000, EXPRESSO) have taken place over the last decade or so. Routine ozone profiling as part of the SHADOZ ozonesonde network takes place at Irene and Nairobi, and aircraft measurements in the MOZAIC programme have contributed both upper tropospheric data and profile data at selected airports. Chemical-transport and trajectory modelling have also increased our understanding of the relative strength of sources and atmospheric dynamics. The purpose of this paper is to synthesize these research results from a variety of sources to present an overview of the key characteristics of tropospheric ozone over Africa.

Attention is drawn, in the first instance, to the importance of Africa as a continent with a large land mass in the tropics and a rich and diverse mix of sources of ozone precursor gases. Highlighted are the contrasts in ozone concentration between east and west Africa and inter-hemispheric differences, the build-up of ozone due to re-circulation in the subtropical anticyclone, the concept of a giant natural photochemical reactor over central and southern Africa and the significance of inter-continental transport to and from the continent

T S Dlamini, CHARACTERISATION OF EMISSIONS FROM SPONTANEOUS COMBUSTION OF COAL AT WITBANK

T S Dlamini 

University of the Witwatersrand, South Africa

Atmospheric pollution associated with coal mining activities results mainly from combustion processes. Spontaneous combustion is one of the most prevalent and serious causes of coal fires. It is a process in which an oxidation reaction occurs without any externally applied heat from a spark, such that temperature rise is due to the coal's heat release through chemical reaction (Querol, et. al., 1998). This incomplete combustion of coal is known to generate poisonous gases (LeRoux, 1972).

Characterizing and quantifying these emissions, which are the main aim of this research, are important steps in air quality monitoring. Gases to be measured are nitrogen oxides, sulphur dioxide, hydrogen sulphide, carbon monoxide, carbon dioxide, ozone and volatile organic compounds. These gases are to be captured using a caravan monitoring station with instruments, within the premises of the Kleinkopje Colliery. Meteorological parameters (wind velocity, wind speed, temperature profile, relative humidity, pressure, rainfall and solar radiation) will also be measured at ground level and this will enable the daily transformations of the pollutants to be determined.

Dietmar Dommenges, A Cautionary Note On The Interpretation Of Eofs

Dietmar Dommenges 

Leibniz-Institut Für Meereswissenschaften, Germany

and

Mojib Latif

EOF and rotated EOF analyses are widely used tools in climate research. In recent years there have been several cases in which the EOF- or rotated EOF analyses were used to identify physical modes. These are the "tropical Atlantic" and the "tropical Indian Ocean SST dipole" modes and the different modes of the Northern Hemisphere winter surface air pressure variability. Here we would like to discuss the problems in interpreting these statistically derived modes as physical modes. By constructing an artificial example we shall show that the patterns derived from EOF- or rotated EOF analysis can lead to misinterpretations. This study should be seen as a cautionary note to highlight the pitfalls which may occur when using the EOF or rotated EOF techniques.

C.B. Du Preez, Classification of the synoptic conditions over the South Western Cape during the ripening period of grapes (February 1996 - 2000)

C.B. Du Preez

ARC-Institute for Soil, Climate and Water, South Africa

and

V.M.F. Bonnardot

V.A. Carey

A preliminary study of the daily weather situations was performed for South Africa and for February months (ripening period of the grapes in the Western Cape), similar to the synoptic classification realized for the temperate latitudes in France (Van Jones & Davis, 2000), in order to study the relationships between climate and viticulture at lower latitudes. Daily weather bulletins of the South African Weather Service (SAWS) and surface data observed at Cape Town International Airport by the SAWS were used. The synoptic weather patterns were classified in four main patterns, namely: the ridging of the Atlantic Ocean High over the western parts of South Africa, the passing of a cold front over the Western Cape, the dominance of the interior trough, and the ridging of the Indian Ocean High over the Eastern parts of South Africa. Of these four classifications, two are applicable in the Western Cape, namely the ridging Atlantic Ocean High and the cold front. The Atlantic Ocean High occurs on 42% of the days in February over the five seasons (1996-2000) used for the classifications, whilst the cold front occurs on 24% of the days. The occurrence of the Atlantic Ocean High varies between 64% (1998) and 32% (1999). Comparing these occurrences with previous research on the influence of vintage (meso-climate) on wine aroma (Carey et al., 2000), it was found, for instance, that warmer conditions in 1998 resulted in predominant tropical fruit aromas in the Sauvignon Blanc wines, and tree fruit aromas in Chardonnay wines. To conclude, it is important to broaden this research to include the whole season because one viticultural stage will influence the others.

Carey V.A., V.M.F. Bonnardot, A. Schmidt & J.C.D. Theron (2000). The interaction between vintage, vineyard site (meso-climate) and wine aroma...South Africa (1996-2000). O.I.V. Bull. (2003) Vol. 76, no863-864., 4-29.

Van Jones G. & R.E. Davis (2000). Climate influences on grapevine phenology, grape composition and wine production and quality for Bordeaux, France. Am. J. Enol. Vitic., Vol. 51. No3., 249-261.

FA Engelbrecht, THE NEW CLIMATE OF SOUTHERN AFRICA

FA Engelbrecht

University of Pretoria, South Africa

and

L RAM

In this paper it is hypothesised that the climate of Southern Africa is steadily settling into a new dynamical mode in response to Global Warming. In the new climate, mid-level high-pressure systems will become well established over the eastern part of the subcontinent in summer. Regions under the direct influence of subsidence underneath these systems (north-eastern South Africa and Zimbabwe) will generally experience drier conditions. In contrast, regions to the north of these highs (Mozambique and Zambia) will experience wetter conditions, since they will be under the influence of stronger easterly winds. Stronger moisture advection around the mid-level highs will cause rainfall to increase over the central to western and southern interior of South Africa. Since these highs will intensify as Global Warming intensifies, the associated cloud bands (and rainfall) are expected to be gradually displaced further and further to the west and south. Upper air troughs from the mid-latitudes will simultaneously be forced to follow more southerly tracks. As the present-day synoptic scale systems settle into their new average positions, certain regions may experience wetter conditions, which will change to drier conditions on a time scale of decades! The changes described above are theorised to be driven by stronger convection in the greenhouse-warmed Tropics. The excess mass that ascends over the Tropics eventually descends over the eastern subcontinent, in this way causing anomalously strong mid-level highs. Winter patterns are hypothesised to undergo similar changes, which are also discussed in the paper.

Simulations of climate change by two high-resolution regional climate models (DARLAM and C-CAM), forced by fully coupled Ocean-Atmosphere General Circulation models (AOGCMs) have been instrumental in formulating this theory. These model results are discussed in detail in the paper.

Acknowledgements

Many thanks to Dr John McGregor, Prof. Hannes Rautenbach and Christien Potgieter for numerous interesting discussions about modelling and climate.

Jesus Fernandez, Precipitation Climatic Prediction Over The Mediterranean Through Statistical Downscaling Of Multi--Model Output From The AMIP2 Project

Jesus Fernandez

Applied Physics II, University Of The Basque Country, Spain

and

Dr Jon Saenz

Dr Francisco Doblas-Reyes

Dr Eduardo Zorita

The Atmospheric Model Intercomparison Project (AMIP) provides a standard experimental protocol for running global atmospheric general circulation models (AGCMs). This kind of project is designed with the goal of assessing the inter--model variability of climate prediction, because the AGCMs are run considering common boundary conditions. Based on 16 models run under the AMIP2 standard protocol, this work estimates the predictability of precipitation over the Mediterranean Basin. The main idea behind this work is to quantitatively evaluate the predictability of precipitation in a region of interest, considering the joint errors derived from the use of state-of-the-art AGCMs and downscaling methods. This may yield an indication of the potential predictability that can be achieved under climate change scenarios. AGCMs do not correctly reproduce precipitation, thus, only large scale fields (namely, SLP) are directly taken from the multi--model output. Precipitation is then obtained by means of statistical downscaling of the model results. Linear (based on Canonical Correlation Analysis) and non-linear (based on analogs) statistical downscaling techniques are used. Since we are concerned with the analysis of second-kind predictability, Probability Density Functions (PDFs) of precipitation for the target region are calculated for the direct output from the models, the downscaled precipitation, the 'mean model' downscaled precipitation and several observed precipitation datasets.

C Ferro, A Survey of Statistical Methods For Climate Extremes

C Ferro 

Climate Analysis Group, Department of Meteorology, The University of Reading, Reading,
United Kingdom

Extreme weather events take many forms and a variety of techniques are useful for determining their statistical properties. I shall focus on methods from the field of extreme-value theory. The survey will review classical methods that are popular in climate research, assess the scope for applying more elaborate methods, and propose areas for future methodological development. Methods will be illustrated using data from the European regional climate modelling experiment PRUDENCE.

Irene Fischer-Bruns, A Modelling Study On The Variability Of Storm Activity On Time Scales Of Decades And Centuries

Irene Fischer-Bruns

Max-Planck-Institute For Meteorology, Germany

The output of several multi-century simulations with a fully coupled OAGCM is examined with respect to the variability of storm activity in winter for both hemispheres on time scales of decades and longer. The frequency of maximum wind speed events within a grid box, using 8 and 10 Bft as threshold, is used as characteristic parameter. Four simulations are considered - two historical climate runs with time-variable solar, volcanic and greenhouse gas forcing of the last five centuries, one control run representative for present day conditions, and a climate change experiment. The main result for the historical simulations and the control run is that storm activity is remarkably stationary. Global maps of storm frequency for different centuries do hardly differ, even though significant temperature anomalies temporarily emerge in these runs. Variations in solar and volcanic forcing as well as in greenhouse gas concentrations within the range of the past few hundred years are not related to variations in storm activity. In the climate change run, which is forced by a 1% increase of CO₂ p.a, however, a gradual change towards more storms in all three storm tracks (North Atlantic, North Pacific, Southern Ocean) is simulated. A more detailed analysis of the variability of storm activity is done with the help of EOFs for both winter hemispheres. The resulting PC indicates the storm activity. The hemispheric mean temperature and the storm activity are decoupled in pre-industrial times. In the transient greenhouse gas scenario, however, the storm activity parallels the development of temperature, and exceeds the two σ -range of pre-industrial variations in the early 21st century.

Maria Dolores Frias, Seasonal Forecast Of Downscaled Maximum Temperature Over The Iberian Peninsula Using Multi-Model Ensemble Versus Single Models By DEMETER Project.

Maria Dolores Frias

University Of Salamanca, Spain

and

Jesus Fernandez

Jon Saenz

Concepcion Rodriguez-Puebla

An assessment is made about the ability of an ensemble of quasi-independent models to produce more reliable regional forecasts than a single model. The multi-model ensemble system developed on the European project known as DEMETER has been considered to analyze the predictability of maximum temperature over the Iberian Peninsula. This project provides seven global atmosphere-ocean coupled models from different European meteorological centers, each running from an ensemble of nine initial conditions. The multi-model ensemble mean and the single models provide large scale information to be translated to the regional scale taking into account the connection between the North Atlantic sea level pressure and maximum temperature over Iberian Peninsula. Statistical downscaling based on Canonical Correlation Analysis is applied in this study in order to estimate maximum temperature in 55 observation sites evenly covering the Iberian Peninsula. Maximum temperature estimated from the multi-model ensemble mean and single models are contrasted with station data by means of correlation, variance fraction and Brier skill score. The results show that the downscaled multi-model ensemble mean underestimates the observed variability, but is more correlated to observations than single models which in general overestimate the variability. Correlation skill is high in early winter and spring and February is the most predictable month.

**T Gill, AN ASSESSMENT OF CHANGING MEAN ANNUAL PRECIPITATION FOR
400 SOUTH AFRICAN RAINFALL STATIONS FROM 1911 TO 2000**

T Gill 

South African Weather Service, South Africa

and

Andries Kruger

The work being presented is a first step in the assessment of changing mean annual precipitation over South Africa. The research is a response to a debate which is currently raging in climate change circles as to whether or not precipitation is increasing or decreasing in the country. A running 30 year normal was calculated on a decadal interval (1911 to 1940, 1921 to 1950 and so on) for each of approximately 400 rainfall stations in South Africa that have sufficiently long daily rainfall time series. No missing data was interpolated and the daily rainfall record that was used was left untouched. The normal values of mean annual precipitation were calculated for each station and trends were determined and mapped. A high degree of variability was found in these trends, both within climate regions and rainfall districts. The trends were then tested for significance and the results mapped. Some explanations are offered for variability which persisted in the significant changes in rainfall, looking specifically at the influence of topography and increasing urbanization, but these studies are far from comprehensive.

The intention in the study is to provide a springboard from which further research can be initiated. Some areas of research that can possibly be undertaken are to apply statistical interpolation techniques to the daily data in order to fill in some of the missing data and then to test the significance of the trends again. The number of precipitation days as well as the intensity of the rainfall could be analysed in order to determine the kind of changes in the rainfall that may be occurring in conjunction with the precipitation trends. It might also be useful to further analyse the trends using stations that have remained fairly rural in nature and compare them to those that have become increasingly surrounded by expanding urban areas.

Nicholas Graham, Ensemble Expansion For Seasonal Climate Prediction Using Recombination

Nicholas Graham 

Hydrologic Research Center, Scripps Institution Of Oceanography, USA

and

Simon Mason

One primary goal of operational seasonal climate prediction systems is to project the probability of relatively extreme seasonal averages or totals. One requirement to make such predictions is accurate knowledge of the probability distribution of the forecast model output. In many cases, the size of the model ensemble may be small compared with the size required to resolve well the probability of a forecast event exceeding some value. We present a method for the expansion of ensemble size for two-tiered seasonal prediction systems (in which the sea surface temperature data is prescribed). The method is based on recombination of individual monthly values from the various ensemble members to construct a much larger population of alternative outcomes. Under the assumption of linear independence of sequential monthly forecast values from any particular ensemble member, these recombined outcomes are equally likely, thus allowing the construction of a large unbiased large population which obeys basic constraints of model physics. An additional benefit is that the relationships between differing variables (for example, temperature and precipitation) is maintained. The results show that if the monthly values are linearly independent, the increase in ensemble size is proportional to the number of ensemble members. We examine the performance of the method using idealized models and with actual AGCM output, and also test the effects of violations of the assumption of linear independence, and of differing monthly variance.

Tomas Halenka, On The Statistical Analysis Of Solar Activity Impact On Stratosphere-Troposphere Global Circulation Patterns

Tomas Halenka 

Charles University, Dept. Of Meteorology And Environment Protection, Czech Republic

The long-term behaviour of global circulation structures and their connections to some heliogeophysical influence, especially solar activity, has been studied for SOLICE Project recently. To represent an objective characteristics of circulation patterns the spectral structure of both stratospheric and tropospheric fields is analysed in terms of spherical harmonics coefficients of expansions for potential vorticity, NCEP/NCAR database of reanalyses is used for period 1948-2002 with monthly data. Temporal analysis of significant spherical harmonics is introduced as well as the comparison of their changes with respect to the changes of different sets of solar, geomagnetic and global circulation indices. A strong connections to a set of extraterrestrial parameters appear for some distinctive shapes of polar vortex as presented in composites for solar maximum and minimum in vertical structure. The natural variability connected to the extraterrestrial influence is studied as well as interannual variability with the emphasis to the QBO and ENSO. The systematic review of the appropriate correlations and linear regression analysis are presented and decadal variability and long-term trends are pointed out for some of wave numbers. Long-term changes in the variability of the circulation patterns are analysed by means of wavelet analysis as well.

Gabriele Hegerl, Detection Of Anthropogenic Climate Change: Using The Past As A Guide To The Future

Gabriele Hegerl 

Division Of Earth And Ocean Sciences, Nicholas School Of Environment, USA

and

Tom Crowley
Francis Zwiers
Peter Stott

Detecting and estimating the anthropogenic contribution to past climate change is necessary to build confidence in predictions of future climate change. Estimates based on the instrumental period will be most complete, but have problems separating the influence of different external influences on climate, such as from solar and greenhouse gas forcing, and do not allow for a well-sampled estimate of the unforced residual noise contribution. Paleo-reconstructions of hemispheric scale surface temperature beyond the instrumental period are more uncertain, but can provide help with both problems. Results are shown using a large ensemble of simulations with a simple climate model to construct a probability density function for climate sensitivity. While these results are important for model validation, society is more concerned about changes in variables that affect it more directly, such as changes in rainfall and climate extremes. The outlook for detection of changes in temperature and precipitation extremes is discussed by studying changes from simulations with two different atmosphere-ocean general circulation models. Results indicate that changes in temperature extremes are significantly different from changes in seasonal temperature, and that changes in extreme precipitation may be more robustly detectable than changes in mean precipitation. Observed changes in annual extremes appear promising, but raise many questions about sampling, homogeneity of time series and difference in trends between different data sets.

Jaakko Helminen, Malting Barley Crop Yield In Southwestern Finland - Potential Risk Management Benefits Based On The Use Of Seasonal Forecasts

Jaakko Helminen

Finnish Meteorological Institute, Finland

and

Ari Rajala

The crop yield and quality of malting barley are largely determined by the temperature and precipitation conditions between the sowing and heading dates. In addition the sowing date, the precipitation between the heading and harvesting dates and the harvesting conditions are of importance. The seasonal forecast deviations standardized with respect to the model climatologies are interpreted as standardized deviations with respect to the observed climatology downscaled to a regular 10 km * 10 km grid by using a combined method of kriging and the Gandin approach, which allows for neighbourhood station choice by relevant physical conditions in terms of geographical information data, like the height, the relative coverage of lakes and the relative coverage of sea with respect to the grid point under processing. By adding to these the relevant threshold values of and risk levels acceptable to the user as well as by considering the costs and benefits related to different decisions made by the user the overall risk management for climate in the context of malting barley cultivation in Southwestern Finland is assessed.

Bruce Hewitson, Cohesive spatial changes of precipitation and seasonality over South Africa

Bruce Hewitson 

CSAG, University of Cape Town, South Africa

While much historical climate change analysis has been undertaken for many regions of the world, this is most often focused on point source data – typically surface station observations. This raises the question over whether the station is spatially representative, and following from this, whether appropriate inferences may be made about change on the regional scale. Compounding this is the question of whether the change is occurring in the base values of the measured variable, or in the statistics or characteristics of occurrence of events.

This talk examines these issues over South Africa using a new gridded data set of daily precipitation, interpolated from a high station density, and with explicit recognition of the variable spatial representivity of each station. This latter attribute is further refined by recognizing that the spatial representivity of any given station is itself a function of the synoptic weather state. The 50-year period from 1950-1999 is examined in terms of the daily precipitation values as expressed in the monthly statistics of wet and dry spell duration, wet-wet and dry-dry day probabilities, median rainfall, number of raindays exceeding a threshold, and the 90th percentile values. The spatial cohesiveness of historical change is assessed in terms of trend, variability, and seasonal changes; collectively showing regional trends within South Africa.

Bruce Hewitson, Empirical Downscaling: Assumptions, Caveats And Stationarity

Bruce Hewitson

CSAG, University of Cape Town, South Africa

Empirical downscaling has, for many impacts researchers, become a staple tool in developing appropriate regional scale climate change scenarios. As regional climate models (RCMs) continue to develop these will perhaps replace some of the empirical approaches; nonetheless empirical tools are likely to remain a key approach for many impacts and adaptation projects. In light of this, and the plethora of approaches present in the literature, it is important to recognize the core assumptions, caveats, and limitations of the conceptual approach (many of which apply equally to RCMs!).

This talk explores some of the foundational issues underlying empirical downscaling, highlighting key areas of concern. In particular the issue of stationarity is considered – an issue commonly put forward as one of the stumbling blocks for climate change applications of empirical downscaling. To conclude, a new downscaling application will be presented, which suggestively shows some convergence between two GCMs for the downscaled future climate over South Africa.

DD Hlatshwayo, The sensitivity of the MM5 model to different parameterisation schemes

DD Hlatshwayo 

South African Weather Services, South Africa

and

CJC Reason

WJ Tennant

Severe Weather is fairly frequent in South Africa and often result in havoc and devastation. Forecasting of severe weather is one of the most important services that is provided by the meteorological profession and is very critical to the protection of life and property. The prediction of severe weather may either be approached by the deterministic numerical weather prediction , the statistical approach or both. For this research the deterministic numerical weather prediction has been used. Model simulations were done using the MM5 mesoscale model nested within the Eta model output.

To investigate the sensitivity of the mesoscale model to the convective schemes in South Africa and to see how the schemes behave for severe and non-severe weather events, two specific convective events are examined. The first case is a severe event that occurred on the 1st January 2002 resulting in a lot of damage over the north-eastern parts of South Africa, and the latter case was a non severe storm that occurred over the east of Pretoria on the 05 February 2004. Two sets of simulations were examined for each case, each with a different cumulus parameterisation scheme on the mother domain. Results show that for both cases, not all schemes in the mother domain function properly with the schemes in the inner domain.

Radan Huth, Detection Of The Modes Of Atmospheric Circulation Variability: Sensitivity To The Principal Component Methodology

Radan Huth 

Institute Of Atmospheric Physics, Czech Republic

We examine the effect of several methodological options in the application of principal component analysis (PCA) to gridded sea level pressure (SLP) anomaly data on the variability modes detected. The methodological options include (i) a compensation for uneven areal distribution of gridpoints in a regular latitude / longitude grid (none, by a cosine weighting or by the use of an equal-area grid), (ii) the selection of similarity matrix (correlation or covariance), and (iii) rotation (none, orthogonal or oblique). The monthly SLP anomalies over the Northern Hemisphere in the winter half-year (November to April) are analyzed. The compensation by a cosine weighting and by the use of the equal-area grid has an almost identical effect if covariance matrix is used. The PCs derived from correlation and covariance matrices differ from each other, the difference being considerably smaller for rotated solutions. Unrotated and rotated solutions differ from each other in the degree of simple structure they possess: much stronger simple structure is present in the rotated PCs. The correlation structure of the data is more similar to the rotated than unrotated patterns. These two facts suggest that in the interpretation of modes of variability, rotated solutions should be preferred. The emphasis is put not only on the leading modes of variability describing the Arctic Oscillation (AO, resulting from unrotated PCA) and North Atlantic Oscillation (NAO, resulting from rotated PCA) features, but also on the higher-order modes which also change their shapes and order of importance in response to the methodological options employed. Concerning the AO vs. NAO dispute, it is important that the Pacific center of the AO turns out to be an effect of high variance rather than of joint variability with its Arctic and Atlantic centers. The study is supported by the Grant Agency of the Academy of Sciences of the Czech Republic, contract IAA3042401.

Radan Huth, The Use Of Multivariate Statistical Methods In The Analyses Of Long-Term Climatic Trends

Radan Huth 

Institute Of Atmospheric Physics, Czech Republic

This contribution introduces two multivariate statistical methods, namely the principal component analysis (PCA) and cluster analysis, into the research on long-term trends in climatic elements, and demonstrates the benefits of their use. The study is based on linear trends of 11 climatic elements at 21 stations in the Czech Republic and is carried out on a seasonal basis. PCA, if applied to the trend magnitudes for a set of variables at a network of stations, allows one to find out mutual relationships among the individual variables, trends. Expected relationships, such as a connection of daily temperature range increases with increases in sunshine duration, and decreases in cloud cover and relative humidity, are confirmed by PCA. In addition to it, PCA allows one to uncover other relationships, which would not have been expected a priori, such as a seasonally varying link between precipitation probability and precipitation amount. PCA can also serve as a tool for identifying the station where the trends are most representative for the whole region analyzed. PCA can be applied to the trends in another way if the data at a single station are stratified, e.g. classified according to large-scale circulation characteristics. Then the trend magnitudes of several variables at a single station under various circulation types enter PCA, which helps to uncover causes of different patterns of climatic change proceeding under different circulation conditions. The cluster analysis of stations allows the spatial consistency of trends to be quantified. The groupings of stations appear to be spatially incoherent and seasonally variable, which indicates that local peculiarities of the relatively complex terrain affect the station trends to a considerable extent. The study is supported by the Grant Agency of the Academy of Sciences of the Czech Republic, contract IAA3017301.

Christopher Jack, Preliminary climate change downscaling with the MM5 regional model

Christopher Jack 

CSAG, University of Cape Town, South Africa

and

Mark Tadross

Bruce Hewitson

GCM based climate change projections are of limited use because of the coarse resolution of the data generated. Grid scales of the order of several hundred kilometers can cover several climate regimes or climatic gradients within one or two grid boxes. Interpreting the data at a human scale is difficult. As a result, much research is being invested in various downscaling techniques. These techniques attempt to add regional scale information to the large scale fields produced by the GCMs. Regional Climate Models (RCMs) are one downscaling technique that can be used. In this paper the PSU/NCAR MM5 regional climate model has been used to downscale three different GCM model datasets. A downscaling of the NCEP reanalysis for the period 1985 to 1995 is presented. A downscaling of the control climate simulation produced by the ECHAM model as well as the future climate (A2 scenario) simulation are also presented. The ECHAM control downscaling is compared to the NCEP downscaling and the differences analyzed both in terms of the differences in the large scale forcing fields and the internal dynamics of the model. Then the ECHAM future and present downscalings are differenced and the differences analyzed in terms of possible climate change signals. Finally a preliminary evaluation of the RCM based downscaling technique is presented in the context of climate change projections. Possible improvements to the technique and future applicability of the technique are discussed.

PA Johnston, Seasonal Forecasts and Agricultural Decision Making - an inside perspective

PA Johnston 

CSAG, University of Cape Town, South Africa

In order to produce more oriented and effective seasonal forecasts it is necessary to gain a greater understanding of the processes and tools involved in the analysis and decision making that a forecast user requires to inform his/her subsequent actions. Several options arise when a user is presented with a forecast - either to alter the current view of things, and consequently his actions, or to ignore the forecast in preference to his own views or for reasons relating to legitimacy, language or method of presentation. There exists for each person a finite amount of information that is used to make decisions. This can be increased through acceptance of new concepts and points of view, but at any given moment decisions will be made using this finite amount of information, influenced by the environment and milieu within which it is made – this is referred to as bounded rationality and it is within this rational space that a decision is made based on the information available. Some forecast users, finding themselves overwhelmed by the array of choice and decision options, settle for the forecast easiest to access, rather than the most suitable.

A greater understanding of a user's frame of reference would be of great assistance to both the producer of the forecaster as well as any subsequent interpreter, facilitator or extension agent. Knowledge of the user's cognitive state leading to decisions concerning the willingness, and thereafter the capacity, to utilise a forecast would provide insight into the degree of importance attached to climate forecast information and the constraints and limitations facing the user in terms of making reactive decisions. The proposed research will investigate maize farmers in South Africa.

Prof Ian Jolliffe, To Centre Or Not To Centre - Or Perhaps Do It Twice?

Prof Ian Jolliffe 

University Of Aberdeen, United Kingdom

The standard definition of covariance or correlation involves centring each variable by subtracting its mean. However, some applications in climatology use uncentred versions. In other disciplines doubly-centred versions have also been employed. A brief review is given of these possibilities, mainly in the context of principal component analysis; their properties are discussed and illustrated on some simple examples.

M R Jury, Spectral modelling of Southern African climate variability

M R Jury 

University of Kwazulu-Natal, South Africa

and

Warren B White
Chris Reason

In this study, statistical techniques are used to decompose southern African summer rainfall into biennial (1.5-3 yr), interannual (3-9 yr) and decadal (>9 yr) bands for the period 1900-1999. With the annual cycle removed, residual spectral energy in the rainfall series is found to cascade from biennial to interannual to decadal bands. Rossby wave action in the South Indian Ocean appears to dominate the biennial scale variability. El Niño-Southern Oscillation (ENSO) and related Indian Ocean dipole patterns are important for interannual variability.

Significant sea temperature and pressure (SST / SLP) fluctuations occurring 6-12 months prior to rainfall contribute biennial and interannual indices to a multi-variate model that demonstrates useful predictive skill. Macro-economic cycles are quasi-decadal, and SST/SLP patterns for this band exhibit eastward wave motion.

Viatcheslav Kharin, Estimating Extremes In Transient Climate Change Simulations

Viatcheslav Kharin

Canadian Centre For Climate Modelling & Analysis, University of Victoria, Canada

and

Francis W. Zwiers

Changes in extreme temperatures and precipitation described in terms of return values of annual extremes are examined in three ensembles of transient climate change simulations performed with the second generation global coupled climate model (CGCM2) of the Canadian Centre for Climate Modelling and Analysis. Three-member ensembles were produced for the time period 1990-2100 using the Intergovernmental Panel on Climate Change IS92a, A2 and B2 emission scenarios. The return values are estimated from a generalized extreme value distribution with time-dependent location and scale parameters fitted to 51-yr samples of annual extremes by the method of maximum likelihood. The method of L-moments for estimating return values is revisited and found to produce biased estimates of return values in the considered transient climate change simulations. The climate response is of similar magnitude in the integrations with the IS92a and A2 emission scenarios but more modest for the B2 scenario. Changes in temperature extremes are largely associated with changes in the location of the distribution of annual extremes without substantial changes in its shape. Exceptions from this general rule occur in regions where land and ocean surface properties change drastically, such as the regions that experience sea-ice and snow cover retreat. Globally averaged changes in warm extremes are comparable to the corresponding changes in annual mean daily maximum temperature while globally averaged cold extremes warm up faster than annual mean daily minimum temperature. There are substantial regional differences in the magnitudes of changes in temperature extremes and the corresponding annual means. Changes in precipitation extremes are due to changes in both the location and scale of the extreme value distribution and exceed substantially the corresponding changes in the annual mean precipitation. Generally speaking, the warmer model climate becomes wetter and hydrologically more variable. Waiting times for precipitation events that are considered extreme at the beginning of the considered simulations are reduced by a factor 1.5-2 by the end of the 21st century.

**Gottfried Kirchengast, Generation Of Global Atmospheric Climatologies From GNSS
Radio Occultation Data**

Gottfried Kirchengast 
IGAM - University of Graz, Austria

and

**Ulrich Foelsch
Andreas Gobiet
Armin Loescher**

The Global Navigation Satellite System (GNSS) Radio Occultation (RO) technique is an active limb sounding method using GNSS radio signals to probe the Earth's atmosphere. Fundamental atmospheric variables such as temperature and geopotential height can be retrieved with high accuracy in the troposphere and stratosphere. The changing thermal structure in this height domain is a sensitive indicator of climate change. The global coverage, all-weather capability, high accuracy, and self-calibrated nature of the RO method suggests it as a promising tool for global monitoring of climatic changes in the atmosphere. The German research satellite CHAMP with a Global Positioning System (GPS) occultation receiver aboard continuously acquires about 180 RO profiles per day since March 2002 and provide the first opportunity to create RO based climatologies on a longer term. The CHAMPCLIM project, a project of the IGAM/University of Graz, Austria, in collaboration with the GFZ Potsdam, Germany, takes this opportunity and aims at exploiting the CHAMP RO data for climate use. After optimizing the RO data processing for climate applications and related data validation, the main emphasis since recently is now to create global RO based climatologies for monitoring climate variability and change on a monthly, seasonal, and annual basis. For this purpose, the complete CHAMP RO data flow (~180 events/day) is used to create refractivity, geopotential height, temperature, and humidity climatologies by two different methods. On the one hand, the climatologies are generated by standard averaging-and-binning techniques, on the other hand, 3D-variational assimilation of the RO refractivity data into ECMWF analysis fields is performed, yielding global climate analyses on a more dense horizontal grid. We will provide an overview on the techniques used and show exemplary climatology results for the JJA 2003 season used as 'testbed' before analyzing the full RO dataset 2002/2004.

Won-Tae Kwon, Comparison Of Extreme Climate Events Between Observations And Simulations Over East Asia

Won-Tae Kwon 

Climate Research Lab, Meteorological Research Institute, Korea

and

Youngeun Choi
Hee-Jeong Baek
Kyung-On Boo

The aim of this study is to understand the trends of the frequency and/or severity of extreme climate events in relation with global climate change. We used the observation data of KMA and the simulation data based on the SRES A2 scenario. Indicators for monitoring changes in climatic extremes worldwide are those recommended by the Statistical and Regional dynamical Downscaling of Extremes for European regions (STARDEX) research project. Using the best available daily data, spatial and temporal aspects of the daily and extreme variables are investigated on an annual and seasonal basis for the periods of 1954-1999 for observation data and for the period of 1951-2100 for simulation data. A systematic increase in the 90th percentile of daily minimum temperatures throughout most of the analyzed areas over Korea has been observed. This increase is accompanied by a similar reduction in the number of frost days and a significant lengthening of the thermal growing season. Although the inter-annual extreme temperature is based on only two observations, it provides a very robust and significant measure of declining extreme temperature variability. The five precipitation-related indicators show no distinct changing patterns for spatial and temporal distribution except for the regional series of maximum consecutive dry days. The regional climate projection over East Asia has been produced using dynamical downscaling nested in ECHAM4/HOPE-G model for 150 years. The model simulation shows about 6°C warming and 10-15% precipitation increase by the year 2100. There are prominent tendency for more drought, more heat wave, less frost day and more heavy rainfall frequencies.

Willem Adolf Landman, Predicting South African Seasonal Rainfall Using A Combination Of MOS And Perfect Prognosis

Willem Adolf Landman 
South African Weather Service, South Africa

and

Lisa Goddard

GCMs typically overestimate rainfall amounts and often spatially distort patterns of rainfall variability. Such systematic biases suggest the need to downscale GCM simulations. Successful recalibration to regional rainfall over southern Africa has been achieved using a perfect prognosis approach and a model output statistics approach (MOS). Here, a method of empirical downscaling is presented where MOS equations are developed using 24-member ensemble GCM simulation rainfall data (the ensemble was forced with simultaneous observed SSTs for each of the 3-month seasons considered) and then 24-member ensemble rainfall real-time forecast fields at different lead-times from the same GCM are subsequently used in these MOS equations. It is therefore assumed that the skill with which the GCM can produce forecast at certain lead-times is as good as skill obtained from simulation data, reminiscent to the assumption of a perfect prognosis approach where “perfect” forecasts are assumed. The ECHAM4.5 predictions are generated 3-month seasons by persisting observed SST anomalies of the month immediately prior to the target season on top of the monthly varying annual cycle of climatological SSTs. At initialization, ensemble members differ from each other by one model day integration for both the simulation and forecast data. Canonical correlation analysis (CCA) is the mathematical technique used to set up the MOS equations and also to do the “perfect prognosis” downscaled forecasts with. These forecasts are verified using ranked probability skill scores (RPSS). It is found that forecast skill is similar to that of the simulations for 0-month lead forecasts of DJF rainfall using observed SST data from November.

C Lennard, THE EFFECT OF NESTING TECHNIQUES IN A REGIONAL CLIMATE MODEL OVER COMPLEX OROGRAPHY

C Lennard 

CSAG, University of Cape Town, South Africa

Orographic features in a given region establish the circulatory patterns of air masses in general. For a mesoscale model to be capable of reproducing this behaviour, a topography is necessary with a resolution such that orographic features are not smoothed away. The simulation domain also has to be large enough to include all the terrain characteristics that are believed to participate in the circulation of air masses, and these can often be far away from the study area. However, modelling such a large domain at such a high resolution makes it computationally impractical. Nesting techniques solve this problem and increase spatial resolution only in the domains where small-scale phenomena might occur and are relevant for the reproduction of all the forcing mechanisms in the study area. At the same time, the outer domain(s) is also included at a coarser resolution, to assure the introduction of larger scale forcings into the inner domain.

The interaction between the domains can be carried out through to different techniques: one-way or two-way nesting. This study examines the effect of different nesting techniques on precipitation in the south-western Cape, specifically around the greater Cape Town region using a regional climate model.

Yun Li, Statistical Modelling Of Extreme Rainfall In Southwest Western Australia

Yun Li

CSIRO Mathematical And Information Sciences, Australia

and

Wenju Cai

Eddy P. Campbell

Rainfall over Southwest Western Australia (SWWA, 118E westward and 32S southward) has decreased over the past decades putting further constraints on water resources in an already dry area. In this study, we analysed daily rainfall over five of geographically dispersed and homogenized weather station within SWWA. We used a peak over threshold method from extreme value theory to model daily rainfall above a given threshold. Change points for extreme daily rainfall were found around 1965 based on different individual stations, with extreme daily rainfall reduced since then. To demonstrate the degree of change in the daily rainfall, we stratified at 1965 and fitted generalized Pareto distributions to the tails of the distributions for daily rainfall in the pre-change period 1930-1965 (including 1965) and the post-change period 1966-2001. The fitted tail distributions also allow the estimation of probabilities of extreme daily rainfall, and quantiles of interest for a given tail area probability. Results show that probabilities and amount of extreme daily rainfall have greatly declined after 1965. There has been vigorous debate as to what forces the drying trend, i.e., whether it is part of multidecadal variability or driven by secular forcings such as increasing atmospheric CO₂ concentration. In this paper, we also use statistical modelling to identify possible associated changes in atmospheric circulation. We find that there is a change point near 1965 in a dominant atmospheric circulation mode of the Antarctic Oscillation (AAO). The result offers qualified support for the argument that the AAO may contribute to the drying trend. This work is supported by a Western Australian government project: the Indian Ocean Climate Initiative (IOCI).

Ta-Hsin Li, An Adaptive Multi-Resolution Method For Estimating Global Temperature Fields From Scattered Observations

Ta-Hsin Li 

Department Of Mathematical Sciences, IBM T. J. Watson Research Center, USA

This paper considers the problem of estimating the entire temperature field for every location on the globe from scattered surface air temperatures observed by a network of weather stations. Classical methods such as spherical harmonics and spherical smoothing splines are not efficient in representing the data that have inherent multiscale structures. This paper presents an estimation method that has the capability of adapting to the multiscale characteristics of the data. The method is based on a spherical wavelet approach recently developed for multiscale representation and analysis of scattered data. Spatially adaptive estimators are obtained by coupling the spherical wavelets with different thresholding (selective reconstruction) techniques. These estimators are compared for their spatial adaptability and extrapolation performance using the surface air temperature data.

**Piero Lionello, Response Of The European Temperature To The Radiative Forcing:
Difference Between Winter And Summer Climate Variability**

Piero Lionello 

Department Of Material Sciences, University Of Lecce, Italy

and

**Simona De Zolt
Juerg Luterbacher
Eduardo Zorita**

This study analyzes the response of the European Climate to the variability of solar irradiance and volcanism during the last five centuries. This period is covered by two different multicentennial global simulations, carried out by the same model and using an identical time dependent RF (which includes, besides the SVRF, also the Green House Gases increase), and by a chronicle-based paleo-reconstruction of the European monthly temperature fields. European temperature is characterized by a higher variability in winter than in summer. In fact, in winter the two simulations present differences of average European temperature that also at multidecadal time scale are much larger than in summer and correspond to different NAO conditions. Spatial distribution of correlation between the two simulation is uniform and it decreases towards North. The correlation between simulations and paleoreconstruction is higher in winter than in summer and has a maximum above North Eastern Europe, while there is no agreement above southern Europe and North Africa. On one hand, these results suggest that predictability of the European climate would be higher for summer than for winter, during which it would be restricted to the multi-decadal or longer timescales at which RF is dominant. On the other hand, they possibly point out to problems in the knowledge of past radiative forcing and climate in the Mediterranean region.

Piero Lionello, Statistical Analysis Of Different Aspects Of Marine Storminess In The Mediterranean Region.

Piero Lionello 

Department Of Material Sciences, University Of Lecce, Italy

and

Antonella Sanna
Alessandro Zardini

This study discusses trends and variability of storminess in the Mediterranean region during the second half of the 20th century. Storminess is evaluated considering three different aspects: monthly synoptic variability patterns, storm tracks, intense surface winds and associate ocean wave fields. Information is based NCAR and ERA-40 reanalysis. The synoptic variability is estimated from the monthly standard deviation maps of the band-pass filtered (1 to 7day cut-off periods) 500 hpa GPH (GeoPotential Height) and SLP (Sea Level Pressure) fields. Storm tracks are identified by an automatic recognition procedure applied to the 6-hourly fields of the reanalysis. The 6-hourly surface wind fields are used for simulating the wave field with the WAM model whose results are considered as a proxy of the presence of intense wind over large areas for a significant duration. The analysis shows the presence of regional processes affecting the storminess in the Mediterranean region beside the well known NAO large scale teleconnection pattern. Moreover, It shows that association to large scale patterns depends on the aspect of storminess considered, so that the regional structures responsible for wave field variability differ from those responsible for the atmospheric synoptic variability. A generalized trend toward reduction of winter storminess has been identified during the second half of the 20th century

Robert Livezey, The Gerrity Equitable Skill Score As A Near Optimal Alternative To The Modified Heidke Score For Verification Of Categorical Seasonal Forecasts

Robert Livezey 

Climate Services Division/OCWWS/NWS/NOAA, USA

The Heidke Skill Score as modified by the NWS Climate Prediction Center and commonly used by other practitioners of climate forecasts has virtually all of the undesirable attributes usually associated with a skill score for ordinal categorical forecasts. These include the ability of forecasters to artificially manipulate the score in several ways to their advantage, inconsistency of the score with other related scores, and no dependence of the degree of forecast error or difficulty on forecaster rewards and penalties. There is an alternative called the Gerrity score that for practical purposes has none of these shortcomings and is easy to compute. The Gerrity score is a particular subset of the Gandin and Murphy family of equitable skill scores for ordinal multi-categorical forecasts. Both the Heidke and Gerrity scores will be described, placed in context with the Gandin and Murphy and other scores available in the literature, and their attributes contrasted.

Robert Livezey, Intraseasonal To Interannual Climate Variability Alphabet Soup: What Is New, What Is The PDO, And Related Questions?

Robert Livezey 

Climate Services Division/OCWWS/NWS/NOAA, USA

In the 1980s a considerable effort (culminating in Barnston and Livezey, 1987) went into diagnosis and cataloging of the principal atmospheric modes of short-term climate variability. The results showed that over North America these modes manifest themselves in the upper-air circulation principally in three patterns or combinations of these patterns, the so-called Pacific/North American pattern (PNA), the North Atlantic Oscillation (NAO), and the Tropical/Northern Hemisphere pattern (TNH). The PNA has been associated with tropical intraseasonals like the Madden-Julian Oscillation (MJO), La Nina, and unforced (by the tropics) internal variability, the NAO with the latter, but the TNH only with El Nino. Since then a number of studies have suggested many other modes or patterns than these, most prominent among them the Arctic Oscillation (AO) and the Pacific Decadal Oscillation (PDO). An obvious addition to the list is the signature (to date) of global change, which is expected to be nonuniform in space and by season (ie not just some uniform warming). A considerable amount of confusion and controversy exists with respect to the AO, the PDO, and global trends. Many argue that for most timescales the AO and NAO are the same phenomena and, for all but very long time scales, are principally a reflection of red-noise internal variability. The growing weight of evidence suggests that the PDO is entirely the reflection of previously-described multiple phenomena, that its characteristic decade to decade reversals have been a consequence of red-noise (thereby unpredictable), and that long-term trends may now be dominating its polarity. As a practical matter, it may be that the only new dimension to the description of short-term (intraseasonal to interdecadal) climate variability is global change, and that as far as wintertime North America is concerned, an adequate description of this variability would encompass MJO, ENSO, global change, and unforced (from the tropics) internal variability. The first three likely have some predictability, the latter none beyond the limits of deterministic atmospheric variability predicted by Lorenzian chaos.

Robert Livezey, Climate Forecasts With Marginal To Moderate Skill And Limitations On Their Value To Real Users

Robert Livezey 

Climate Services Division/OCWWS/NWS/NOAA, USA

and

Barbara Mayes

Marina Timofeyeva

Long-lead seasonal forecasts of the 1997-98 El Nino and the 1999-2001 La Nina and their impacts (especially for cold season North America) were made with unprecedented skill, detail, and confidence. This success has led to enormous interest in seasonal prediction worldwide, including in developing countries, and in many instances unrealistic expectations about them. The fact remains that climate predictions have at best modest skill, and in many circumstances no or marginal skill, in the absence of a strong ENSO signal. Nevertheless, it has been argued forcefully by a number of groups that these sets of forecasts have tangible economic value for a class of decision-makers and users. The arguments are based on application of simple cost-loss models, which relate forecast value in a simple decision environment to various measures of forecast performance and skill. The credibility of the cost-loss arguments depends on whether real decision-makers and users behave in the manner assumed in the models. In existing applications of the models, the value of the forecasts is the mean expected over many consistent iterations of the forecast/decision process. In the case of a winter forecast, a particular user can only exercise this process once per year a particular location. Thus, for the cost-loss model value estimate to be relevant to the user, he/she must exercise the process over many years and/or over many independent locations in a disciplined, consistent manner. It is difficult to imagine a user that would do this over a period of even a decade, or whose decision-making environment would remain static for that long. Clearly a potential user needs to understand (1) his/her cost/loss situation, (2) the expected skill of the forecasts for those parameters, places, and times of year of interest to them, i.e. on which their decisions will be based, and (3) what is necessary to ensure a high likelihood of realizing value from use of the forecasts. Here we will address (2) and (3) in the context of official U.S. long-lead seasonal temperature and precipitation forecasts issued by NOAA's Climate Prediction Center. We begin by computing and combining skills in ways that are informative to potential customers of the forecasts. We first display the bulk skills by lead time (which in itself turns out to be informative) and then provide the same graphs stratified by time of year, major ENSO episodes, and both. Where possible we combine consecutive leads with similar skills to achieve more confident estimates. For those cases that warrant further consideration we stratify by large-scale U.S. regions (like east vs. central and west or south vs. north). We demonstrate that some forecast subsets have no or marginal skill, while others have moderate or even substantial skill. Next we present estimates of the uncertainty in forecast value for different forecast skills and realistic iterations of the forecasts/decisions, as well as with different simple assumptions about the psychology of the user (for example, the user abandons use of the forecast if the forecast was wrong two winters in a row). The computed uncertainties cast considerable doubt on the utility of marginally skillful forecasts sets for individual users and provide a sense of what skill levels are necessary to increase likelihoods that relatively short sequences of forecasts/decisions will be of value. These skill levels turn out to be relatively high, comparable to the performance of the strong ENSO event "forecasts of opportunity." On the other hand, users who can exercise forecast/decisions at multiple independent locations each season over two or more seasons a year (like weather derivatives brokers) can be reasonably confident of realizing value from only moderately skillful forecast sets.

Robert Lund, Trends In United States Maximum And Minimum Temperatures

Robert Lund

Department Of Statistics, The University Of Georgia, USA

This talk studies linear trend rates in high and low temperatures recorded in the contiguous 48 United States over the last 150 years. Trends computed from extreme value and regression based methods will be compared. Methods for handling station location moves at known times (change points) will be discussed. Spatial contour maps of the annual warming/cooling rates were constructed via the nonparametric head-banging algorithm; these rates are partitioned into seasonal trends for Spring, Summer, Fall, and Winter.

Neil MacKellar, REPRESENTING THE LAND SURFACE IN A REGIONAL CLIMATE MODEL: WHAT ARE THE OPTIONS

Neil MacKellar 

CSAG, University of Cape Town, South Africa

and

Mr Mark Tadross
Mr Bruce Hewitson

Numerous sensitivity studies have confirmed the importance of land-surface boundary conditions in atmospheric modeling. It can therefore be inferred that, in order to work towards increasing the accuracy of weather and climate simulations, it is essential to consider the quality of land-surface data used by the model. Two possible sources for such data are satellite observations and dynamic global vegetation models, each with its respective attractions and caveats. In this study, the nature of some land-surface data sets available for southern Africa are described, and the practicalities of how this information may be incorporated into Version 5 of the Penn State/NCAR Mesoscale Model (MM5) are explored. In addition, the Sheffield Dynamic Global Vegetation Model (SDGVM) is employed to simulate a vegetation map for use by MM5. The sensitivity of MM5 to using this simulated land surface as opposed to the default configuration is assessed for a 3-month summer period. This represents a first step in investigating the possibilities for a dynamic coupling of these two models.

A Manhique, Atmospheric and ocean patterns associated with rainfall anomalies in Mozambique

A Manhique 

Department Of Oceanography, University of Cape Town, South Africa

and

**C.J.C. Reason
Y Richard
Margaret Rouault
L Rydberg**

Atmospheric and sea surface temperature (SST) patterns associated with rainfall anomalies in Mozambique are studied. The study analyses the spatial and temporal variability of rainfall using extended empirical orthogonal functions (EEOF). Southern Mozambique appears to have strong spatial and temporal coherence in the late summer (January – March (JFM)). Some coherence is also observed in the early summer (November – December (ND)) in the north. Correlation analyses applied between successive months confirm a strong – positive correlation in the southern region. However in the north, the correlations are weak. Nevertheless two regions (south: 20 - 26 S; north 10 - 15 S) and three periods (October; ND; JFM) were considered and respective indexes created. The indices were created by averaging the standardized monthly rainfall anomalies for each region.

Composite analyses show that positive rainfall anomalies in the north region during early summer (ND) are associated with cooling in the north and equatorial Indian Ocean. Simultaneously, some increase in SST is observed south of Madagascar. Atmospheric patterns suggest shifts of the intertropical convergence zone (ITCZ) associated with northeast monsoons as the main mechanism driving the rainfall anomalies. When the ITCZ is positioned further south, the region tends to receive more rain, and less when it is located more to the north. The JFM rainfall anomalies in the south region appeared strongly connected to global patterns with shifts in the Walker circulation being important.

Lizelle Maritz, Spatial And Temporal Variation In Rainfall In The Arid Zone: Is It Changing?

Lizelle Maritz

ARC - Institute For Soil, Climate And Water, South Africa

and

Johan Malherbe

Anneke Thackrah

Frans Koch

In general, arid regions tend to have a higher precipitation variability, both spatially and temporally, compared to humid regions. To examine this variability two sets of rainfall surfaces were used: Seasonal rainfall surfaces (July to June), using climatic data spanning from 1930 to 2003, and dekadal rainfall surfaces, covering a 13-year period (1986 - 2001, excluding 1994 and 2000). Both these sets of surfaces were developed by the ARC-ISCW using weather station (SAWS and ARC-ISCW) and satellite data (for dekadal rainfall surfaces). The Köppen classification criteria were used to define the arid zone that was used as the study area. Köppen divides the arid zone into annual, summer and winter rainfall areas thereby already stressing the spatial variability of rainfall in the zone. Rainfall surfaces and point data (weather stations reporting between 52 and 101 years) were analysed to ascertain whether any significant spatial and temporal changes occurred within the study area both inter- and intra- annually and seasonally.

Ruth Thokozile Massey, Climate Change, Related Risks And The Role Of Non-Climate Related Stressors: Perceptions, Adaptations And Responses Of Rural Households In Kwa-Zulu Natal

Ruth Thokozile Massey 

University Of The Witwatersrand, South Africa

Climate related risks within Southern Africa threaten the livelihoods of many communities particularly those who rely on a variety of resources for their subsistence. This risk is primarily determined by the situation or context in which communities find themselves. Adaptation to and coping with climate variability are key themes in current discussions and policy initiatives. The Thukela Catchment region in rural Kwa-Zulu Natal, South Africa, was used as a case study for this research. Two communities were compared in the study. The first has the Mudane Irrigation Scheme (MIS) organised mainly by the local Department of Agriculture (DoA). The second has no formal irrigation scheme and is relatively untouched, directly, by formal institutions. The region is currently in a drought situation but has had recent flooding. The primary aim of this research was to determine and better understand how the two communities perceive and respond to climate-related risks and to look at the role non-climate related stressors have on their ability to cope with climate variability and environmental change. Focus was placed, specifically, on adaptation and coping and on why some households within the same community cope with climate related stress better than others. The role of formal institutions and food security were looked at in particular. Findings have shown that the first community relies heavily on DoA for access to information, training and education as well as access to water, physical capital, infrastructure and decision making arenas. This in fact increases their vulnerability and restricts their ability to cope and adapt. This paper raises questions around the role of adaptation and formal institutions in the livelihoods and coping strategies of households with regard to climate change.

Andrew Matthews, Understanding Drought: A Four-Dimensional Generalized Linear Model Of Sahel Rainfall

Andrew Matthews 

Climatic Research Unit, University of East Anglia, United Kingdom

and

Jean Palutikof

Tom Holt

A generalized linear model is used to link rainfall in the Sahel to the leading modes of atmospheric variability over Africa and Asia. The model is an attempt to identify the lagged atmospheric processes underlying the persistent dry conditions over the region since the late 1960s. Daily rainfall data from over 500 stations in sub-tropical western/central Africa were gridded to a 1° resolution. Principal Component Analysis (PCA) was used to group the gridded rainfall data into six spatially coherent areas, forming the predictand regions in the regression experiment. The main modes of atmospheric variability over a domain stretching from the mid-Atlantic to Indonesia, at four atmospheric levels, were identified using a four-dimensional PCA of six variables extracted from the NCEP Reanalysis climate data set. The lagged scores from the PCA are the predictors in the regression. The generalized linear model linking the lagged atmospheric predictors to the rainfall indices accounted for up to 30% of the rainfall variability, providing a good representation of the gamma-distributed rainfall series.

T N Mdluli, The Societal Dimensions of Domestic Coal Combustion and Alternative Energy Sources

T N Mdluli 

University of the Witwatersrand, South Africa

and

Luanne Otter
Coleen Vogel

Air pollution is one of many issues that have a direct impact upon the economy and the well being of society in South Africa. Domestic coal combustion contributes significantly to the air pollution problem in the country. Mitigation measures that can be put in place to reduce emissions from domestic coal combustion may have a direct impact upon the environment and the societies that burn coal. The aim of this research, therefore, is to provide an assessment of the societal impacts of mitigating emissions from domestic coal combustion in Soweto. A questionnaire survey will be conducted and it will explore fuel-use and socio-economic data, health impacts of domestic coal combustion and the society's acceptance and willingness to pay for measures that reduce air pollution.

J.-L. Melice, Climate variability at sub-Antarctic Marion Island since 1960

J.-L. Melice 

IRD, Laboratoire d'Océanographie Dynamique et de Climatologie, France

and

Mathieu Rouault

Chris Reason

Johann Lutjeharms

Marion Island (47S, 38E) climate has dramatically changed since 1960. During the last 40 years, a significant increase of the pressure and of the air and sea surface temperatures is observed together with a significant decrease of the rainfall and a northward shift of the wind direction. We show that the trend in the pressure observed at Marion is linked to a change in phase and amplitude of the Semi Annual Oscillation (SAO) index and could explain the trends observed in the temperature, wind direction and rainfall. The changes observed in the SOA index can be linked to recent trends in the lower stratospheric polar vortex which, according to Thomson and Solomon (2002), are due largely to ozone loss.

REFERENCE

Thomson, D.W.J. and S. Solomon (2002). Interpretation of recent Southern Hemisphere climate change. *Science*. 296, 895-899.

Ladislav Metelka, Non-Linear SLP Variability In North Atlantic Region In Winter Period

Ladislav Metelka 

Czech Hydrometeorological Institute, Czech Republic

The North Atlantic Oscillation (NAO) is the most significant oscillatory system in North hemisphere extratropics. It can be simply described with the help of standardized anomalies of SLP differences between one station in Icelandic region and the other in the region of Azores or in SW Europe. More sophisticated approach to the NAO is based on Principal Component Analysis (PCA) of SLP or geopotential heights at low tropospheric levels in North Atlantic region. As PCA is linear method, the results of PCA are influenced by the application of linear method to the non-linear problem (interpretation of the results of non-rotated vs. rotated PCA, etc.). Neural networks belong to Artificial Intelligence (AI) systems. One of the special kinds of neural networks, so called “autoassociative neural network” is able to perform the non-linear counterpart of PCA. This method has been used in our study for the non-linear description of winter North Atlantic SLP variability. Due to the fact that the original SLP values in nearby grid points are highly intercorrelated, linear PCA has been used for data pre-processing (for dimensionality reduction only, not for description of the individual oscillatory systems). Then the autoassociative neural network non-linearly combining several linear PCA scores to a single non-linear score has been built. Results of our study indicate that the SLP variability in North Atlantic region is non-linear in its origin. It is indicated by:

- the shift between positions of NAO centers in the positive and negative NAO phases
- some non-linear links between NAO and other NH oscillatory systems (especially Scandinavian Oscillation) which can not be revealed with the help of linear methods
- signs of “rotational exchange” of the individual centers of SLP anomalies during the transition between positive and negative NAO phase.

P.S Moatshe, Verification of South African Weather Service operational seasonal forecasts.

P.S Moatshe 

South African Weather Service, South Africa

and

Willem Landman,

The value of evaluating forecast accuracy from the user's perspective is important. Through understanding of forecast performance, it helps decision makers to determine when and how to respond to expected climate anomalies. Evaluation of seasonal forecasters also can help planners reduce their vulnerability to climate, because they can plan more informed and prepare more effectively. A probability forecast can be evaluated using different skill scores, but in this case the Ranked Probability Skill Score (RPSS) is used. RPSS is the Ranked Probability Score (RPS) of the forecast compared with the RPS of the forecast of climatology that assigns 33.3% for each three equi-probable categories. The RPSS measures the cumulative error between the categorical forecast probabilities and the observed category relative to some reference forecast. If the dominant forecast category was observed, the RPSS is positive (performing) and if not the RPSS is negative (under-performing). RPSS is zero when there is no skill.

The SAWS issues probabilistic seasonal forecasts every month. These forecasts are a result of expert interpretation of output from a large variety of seasonal forecast models. Verification of five January to March (JFM) season is presented at different lead-times ranging from zero to four months.

G V Motsa, Atmospheric sulphur transport over the South African Highveld.

G V Motsa 

Climatology Research Group, University of the Witwatersrand, South Africa

and

Stuart Piketh

Industries located on the South African Highveld introduce a wide range of anthropogenic primary pollutants directly into the atmosphere, including particulates such as ash, soot and dust ; gases such as nitrogen dioxide, nitrous oxide and sulphur dioxide. The industrial Highveld region accounts for 90% of South Africa's scheduled emissions. Secondary pollutants, such as nitric acid, sulphuric acid and sulphates are formed over time from primary pollutants through chemical conversions and oxidation in the atmosphere.

This research will focus on sulphur dioxide concentrations in the atmosphere over the Highveld and how these concentrations change with altitude and time. The main aim is to demonstrate trends in sulphur transport over the Highveld. Peak concentrations of sulphur dioxide in the atmosphere will be established and their exact locations -especially in relation to sources on the Highveld - will be determined. The rate of decrease in sulphur dioxide concentrations in plumes over space and time will also be determined.

The preliminary results indicate that sulphur dioxide concentrations over the Highveld are above background levels especially over and near sources. Highest sulphur dioxide concentrations occur at altitudes between 1500-2000m. These preliminary findings indicate tall stack emissions as the main source of sulphur dioxide in the atmosphere.

NS Mpandeli, Use And Value of Seasonal Climate Forecasts in Rabali Area

NS Mpandeli 

Agriculture Research Council - Institute for Soil, Climate and Water, South Africa

and

MT Bila

T Ramudzwagi

K Mahuluhulu

E Archer

The IPCC and the climate-impacts modeling work undertaken at IIASA, amongst others, point to the possibility of the southern African region being negatively impacted by climate variability and change. Much progress has been made in atmospheric modeling work, including general circulation models and seasonal forecasts although much still needs to be done. Despite the lack of certainty in the use of such models, there is sufficient progress that has been made to begin considering how such products could possibly be used by agriculturalists in their decision-making processes. With this as backdrop, the use and value of seasonal forecasts as a potential agricultural risk-management tool by farmers in the Northern Province, Rabali area is examined. This paper furthers the debate on the user uptake of climatological and meteorological products in the southern African region. Several questions are addressed including: When do farmers normally require such information? How is such information currently collated and used? How are forecasts currently perceived and how could this perception, if negative, be improved?

SK Mulaudzi, An estimate of emissions from domestic biofuel combustion in Southern Africa.

SK Mulaudzi 

University of Witwatersrand, South Africa

Combustion of fuelwood, charcoal and non-woody biofuels is a daily practice for about half of the world's population. Combustion of biofuel is a major source of trace gases, with domestic biomass burning contributing about 17% carbon dioxide (CO₂), 13% carbon monoxide (CO) and 6% nitrous oxide (N₂O) to the global budget. In Africa, where there is a growing population, domestic biofuel emissions are particularly an important source of trace gases. The most important source of biomass fuels in Africa is wood fuel (Fuel wood and charcoal), crop residues and animal dung. In this project, the amount of fuel (wood, dung or crop residues) used in rural households across Southern Africa is measured daily over winter and summer months. This data is used to estimate the trace gas production from domestic fires throughout Southern Africa. The research also compares the trace gas emissions between different Southern African countries and discusses the seasonal changes in emissions.

T Ndarana, The performance of the Eta model when its data assimilation is removed

T Ndarana

South African Weather Service, South Africa

and

Hilarie Riphagen

Warren Tennant

The South African Weather Service is currently running the fourth version of the operational Eta model with a 32 kilometre horizontal resolution and 45 vertical Eta levels. The model has its own data assimilation procedure which is a twelve hour process and takes about 22 minutes (15.7% of the total run time) to run on the Cray SV1. The procedure involves the use of twelve hour old global fields from the NCEP global spectral model on the Global Telecommunication System (GTS) as initial guess and boundary conditions, three dimensional variational analysis and four three-hour incremental runs.

The aim of this study is to test the model performance when the data assimilation process is removed. To obtain the model initial conditions, the latest GTS fields which contain observations that cover the whole spectrum of observation types are employed. These are made available at 09h00 and 21h00 (GMT) on a daily basis.

The model accuracy, skill and reliability for both the operational and the experimental model runs are evaluated and compared. This is done for both 24 and 48 hour forecasts. The evolution of the skill of the different models investigated. The statistics of each system are evaluated against each model own analyses.

Mark New, Estimating Oceanic Precipitation fields in the Pre-Satellite era

Mark New 

Geography Department, Oxford University, United Kingdom

and

Dimitrios Efthymiades

Richard Washington

A fundamental limitation of gridded precipitation datasets is the paucity of observations over the oceans in the pre-satellite era. We have evaluated the capabilities and limitations of reconstructing seasonal oceanic precipitation, using land-based precipitation records. The reconstruction method is based on eigenvector projection, where eigenvectors (or 'EOFs') of precipitation variability in the CMAP merged, gauge and satellite-based, precipitation dataset, with near-global coverage from 1979-present, are projected onto land-only data. The approach has considerable skill in parts of the Tropics where interannual variability is mainly associated with ENSO activity. In non-ENSO years (for the tropics) and for all years in most of the extratropical oceans, reconstruction skill is substantially reduced in areas more than 1000km from land data points.

Mark New, Probabilistic Regional Climate Prediction: Potential And Pitfalls

Mark New 

Geography Department, Oxford University, United Kingdom

In the last few years simple and intermediate-complexity climate models have been used in Monte Carlo experiments to provide estimates of the probability distribution of future global mean temperature. While these results are useful for quantifying the limits to our present ability to predict future climate on global scales, they provide little information of relevance at the regional and local scales at which impacts and adaptation options are assessed. New developments in Monte Carlo simulations of coupled AOGCMs will in the next few years offer the potential for probabilistic predictions at regional scales and, with appropriate downscaling, at local (sub GCM-resolution) scales. Provision of limits and/or probability distributions for future climate at these scales offers impacts and adaptation researches a powerful opportunity to assess options in a risk-based framework. However, these *conditional* probability distributions for future climate and its impacts are much easier to misinterpret than scenario-based approaches, as they provide the end-user with the illusion that they are unconditional probabilities.

L. D. Ntsangwane, Investigating the optimal use of a super ensemble over southern Africa

L. D. Ntsangwane 

South African Weather Service, South Africa

and

Willem Landman

Richard Sewell

The ensemble rainfall simulations for the 29-year period for the December to February (DJF) season from five General Circulation Models (GCMs) are considered to investigate the optimal use of a super ensemble over Southern Africa. The number of ensemble members from each GCM varies from 9 to 24 with 63 ensemble members in total. One possible approach of incorporating model sensitivities into ensemble forecasting systems is to combine ensembles run from two or more GCMs. The statistical approach known as the Kolmogorov-Smirnov test is used to test the hypothesis that the simulations PDF from the individual ensemble systems are taken from the same population. The results of the five GCM ensembles show that, at 90% level of significance all the GCMs are somehow significantly different. Therefore, all the GCM ensembles are combined to propose an effective ensemble size. This will however, lead to probabilistic forecasts that are skilful over a greater portion of the region and a greater portion of the time series. The Rank Probability Skill Score (RPSS) is used to calculate the effective ensemble size of a super ensemble suite. In addition, the results indicate that even after combining all the GCM ensembles, not all the 63 ensemble members of a super ensemble are required for optimal skill.

Heiko Paeth, A Dynamical-Statistical Approach For Seasonal Forecast Of Subsahelian Rainfall Using Cross Validated Model Output Statistics

Heiko Paeth 

Meteorological Institute, University Of Bonn, Germany

and

Andreas Hense

Rainfall over sub-Saharan West Africa is subject to large interannual variations, strongly affecting agricultural planning. This study deals with the predictability of interannual rainfall anomalies during the rainy season (June-September) over sub-Saharan West Africa. First, regional and global near-surface predictors for sub-Saharan precipitation are derived from multi-model ensemble output. Second, a cross validated stepwise multiple regression model is developed, choosing physically motivated predictors. This statistical model is used as model output statistics system (MOS) to forecast the rainy season. This attempt goes one step further than eg. the Krishnamurti approach by deriving observed rainfall anomalies from simulated variables other than rainfall in order to avoid systematic errors and uncertainties in the modelled convection, cloud and rainfall processes. The forecast skill of the MOS system is found to be small but significantly larger than zero. A low-cost method for operational seasonal forecast is proposed. Sub-saharan rainy seasonal precipitation (RSP) is closely connected with tropical Atlantic sea surface temperature (SST). However, there is also a significant teleconnection to El Niño and La Niña events in the eastern equatorial Pacific. In addition, RSP reveals a strong negative correlation with sea level pressure (SLP) over the tropical Atlantic and the West African subcontinent, indicating the West African monsoonal strength. The most important predictors for observed precipitation are represented by the SLP variables, regional mean and gradient, rather than the simulated rainfall itself. Compared with the null-hypothesis based on randomized predictor chronology, the multi-model predictability is statistically significant. The validation reveals a significant forecast skill, especially with respect to small rainfall anomalies. Given the distinct autocorrelation of tropical SST, this study holds the prospect of operational seasonal forecast over a broad sub-Saharan region with low-cost methods.

Heiko Paeth, A Bayesian Decision Method For Climate Change Analysis With Two Application: Temperature And NH Circulation

Heiko Paeth 

Meteorological Institute, University Of Bonn, Germany

and

Seung-Ki Min
Andreas Hense

Climate change analysis can also be viewed as a decision or selection process of the scenario - greenhouse gases (GHGs), GHG plus sulfate aerosols (SUL), or natural variability - which is most probable in view of the observations. This decision can be made by means of the Bayesian decision theory. Based on NCEP reanalysis data and model data from multi-model ensembles under control and GHG(+SUL) conditions, it is questioned whether the observed climate characteristics during the 20th century are classified in the undisturbed case or one of the two changed climates. It is also addressed to which extent the Bayesian decision results are sensitive to the choice of the prior probability for the three climate scenarios and the estimate of natural variability. Two application studies have been made: (1) A two-dimensional approach refers to the lower tropospheric and stratospheric temperature. (2) A one-dimensional approach uses the index time series of the Arctic Oscillation (AO). In terms of temperature, it is found that the observed anomalies from the late 1990s onward stand out from natural variability, even if small prior probability is assigned to the disturbed climates. The observed AO index anomalies during the 1990s are also classified in the simulated GHG scenario, but recently the signal drops down. Instead, the long-term AO trends reveal a clear climate change signal, which is robust against the choice of the prior. The estimate of the natural variability is a crucial factor in the Bayesian decision theory, but does barely affect the results in terms of the AO.

Jan Picek, Tail Index Of Maximum Air Temperatures And Precipitation In Czech Republic

Jan Picek

Technical University In Liberec, Czech Republic

If we are interested in the extremal events such as the extreme intensity of the wind, the high flood levels of the rivers or extreme values of environmental indicators then we are rather interested in the tails of the underlying distribution than in its central part. A goodness-of-fit test usually concerns the central part, hence it does not provide us with a sufficient information on the shape of tails. Our primary goal is to study more closely the shape of the tail. The authors recently constructed several nonparametric tests of one-sided hypotheses on the value of the Pareto-type tail index of the innovation distribution in the linear autoregressive model. The tests are nonparametric and are based on the series of residuals with respect to an appropriate estimator of the AR parameters; more precisely, they are based on the empirical process of maximal residuals of non-overlapping segments of such series. Inverting the tests in the Hodges-Lehmann manner, we obtain strongly consistent estimators. The estimators are applied to the series of the maximal daily temperatures and precipitation at the Czech Republic station in the period 1961-2000.

Rudi Pretorius, Uncertainty, Scientific Credibility And Related Issues Concerning Global Climate Change

Rudi Pretorius

University Of South Africa, South Africa

Uncertainty, scientific credibility and related issues concerning global climate change The contribution by WG1 to the TAR presents new and stronger evidence that most of the warming over the 20th century (in total $0.6 \pm 0.2^{\circ}\text{C}$) is attributable to human activities and that the human influence will continue to change atmospheric composition throughout the 21st century. It has to be noted, however, that the extent to which it is possible to distinguish between the anthropogenic influence and the background of natural climate variability, remains debatable - despite the progress which has been made to reduce uncertainty. In addition, projections of the climate of the future, which are to a large extent supplied by global climate models with their own limitations and uncertainties, are inherently not very specific either. An example is the projection supplied in the TAR of a global average surface warming from 1990 to 2100 ranging from 1.4 to 5.8°C , based on a number of climate models and on the full range of emission scenarios provided in the SRES. Compared with the earlier IS92 range of 1.0 to 3.5°C as reported in the SAR, the TAR range is significantly greater, with the upper-end significantly higher as well. This paper presents a critical analysis of the latter as well as other findings of the IPCC in its assessment reports (with focus on the TAR) concerning climate change science and also reviews and analyses the relevant debates and criticisms in this regard. The extent to which anthropogenic factors are responsible for observed climate changes is considered, with reference to the associated disagreements. These can largely be ascribed to the numerous uncertainties about global climate change which still exist. In the light of these uncertainties, together with the fact that the complexity and unpredictability of the earth's climate system and the interactions among its components are apparently still underestimated, this paper argues that scientists should acknowledge that accurate projections of the future climate together with its impacts are still not possible at this stage. Considering the lack of sufficient hard evidence, and in order not to lose scientific credibility, this paper concludes that it is crucial to play open cards and to distinguish clearly between the 'facts' and the 'fiction' regarding global climate change.

Nazario Ramirez, A Transfer Function Model And Remote Sensing Data To Estimate Soil Moisture

Nazario Ramirez 

Department Of Industrial Engineering, University Of Puerto Rico, Puerto Rico

and

Ramon Vasquez

Harold Cruzado

It is well established that performances of numerical models are very sensitive to initial and boundary conditions. The regional atmospheric modelling system (RAMS) that is used to simulate with high resolution the climate dynamics in Puerto Rico is highly sensitive to soil moisture initial conditions. Incorrect initial conditions will generate misleading modelling results. Therefore, this research effort is focused on the development of reliable estimates of soil moisture. Remote sensing measurements, artificial neural networks and a transfer function model are used to estimate soil moisture process in Puerto Rico on a daily and monthly basis. Preliminary results of an on-going project will be presented in this research work. The components of water balance system are: precipitation, runoff, evapotranspiration, and groundwater loss. The driver of the soil moisture system is the amount of precipitation that occurs in a given area and time. The runoff is highly dependent on the rainfall intensity, roughness, vegetation and soil type. The evapotranspiration is mostly associated to rainfall intensity, air temperature, and wind velocity. The amount of percolation will also depend on rainfall intensity, soil, and vegetation type. The major assumption of this research establishes that the soil moisture is a function of the following observable variables: precipitation, air temperature, elevation, soil type, and vegetation index. Therefore, a transfer function (TF) model is suggested to represent the interactions of the variables involved in the soil moisture process. The proposed model pretends to represent the spatial and temporal variability. The spatial variability will be modelled by using a self organized neural network (NN) including the following input patterns: vegetation index, rainfall, air temperature, soil type, and elevation. The spatial variability will provide a reasonable number of homogenous regions and inside of each region a temporal variability will be model by using the transfer function model. The input variables of the TF model are: rainfall, air temperature and the output of the TF model is the soil moisture. The TF model represents the stochastic behaviour of the soil moisture for a specific soil type, vegetation index, and elevation. Thus, the outputs of the TF model will provide estimation of soil moisture taking into account the spatial and temporal variability. The outputs of the transfer function will be daily and monthly maps at 1 km² of resolution of Puerto Rico soil moisture. Satellite data is obtained from NASA Goddard Space Flight Center at the Distributed Active Archive Center (DAAC). The MODIS sensor provides air temperature and vegetation index for Puerto Rico at 1 km² of resolution. Two passes of instantaneous air temperature are given on a daily basis. One pass represents about the maximum temperature and the other the minimum temperature; since the first pass occurs about noon and the second one during the night. Since MODIS does not provide air temperature when clouds are present, a second NN is designed to interpolate air temperature under the presence of clouds. The Backpropagation algorithm with the Levenberg-Marquardt (LM) learning rule was used to accomplish this interpolation task. The LM learning rule was used to accelerate the convergence of the NN. The input patterns of the NN are: elevation, vegetation index and rainfall. The target of the NN was the air temperature obtained from MODIS and

from 52 weather stations. The daily cumulated rainfall for Puerto Rico was obtained from radar data at 5 km² of resolution. The air temperature that comes from stations was provided by the National Climate Data Center and the daily rainfall data was provided by the National Weather Service. Elevation and soil type were provided by the Laboratory of Remote Sensing from the University of Puerto Rico.

CJ Rautenbach, LIKELY FUTURE CHANGES IN TEMPERATURE AND RAINFALL IN THE KOMATI RIVER CATCHMENT BASIN

CJ Rautenbach

University of Pretoria, South Africa

and

**Stuart Piketh
Jonas Mphepya**

Climate change and variability has received a large amount of attention over the past three decades. Concerns about rapidly changing global surface temperatures in response to elevated concentrations of greenhouse trace gases such as Carbon dioxide have legitimately been raised. In 2002 the International Panel on Climate Change (IPCC) issued a series of studies that evaluated the scientific basis for climate change as well as the most likely problems that may result and the ability of nations around the world to adapt to these changes. Southern Africa was identified as one of the regions of the world that would not adapt easily to changing climates, in particular increases in extreme events, mostly as a result of a lack of resources. Given the aforementioned it is important for southern African nations to start to plan for the possible impacts that may arise from increases and or decreases in such climatic factors as temperature and rainfall. With this in mind Eskom has started to evaluate the implications of possible climate change to current and future power utility operations and locations. The research has two primary objectives that are pursued as a measure of precaution to prepare for likely changes in future climates:

- To determine whether any IPCC General Circulation Model (GCM) inter-model consistency appears in future anthropogenic climate change projections over South Africa with specific emphasis to the Komati river catchment.
- To identify any significant trends in more recently observed climate patterns that concur with consistent IPCC GCM future climate change projections.

The research has been subdivided into three main activities. The first is to study climate change simulations produced by six of the most advanced GCM's in the world. Inter-model consistency increases the validity of the predictions. Predictions over the next fifty years are considered. The second component attempts to evaluate the changes within the Komati river catchment. This has been done by nesting a higher resolution model (DARLAM) within the larger scale CSIRO GCM. Last, an attempt is made to use observational data from the Komati River catchment from surface weather stations to evaluate if any changes in surface temperature, rainfall and mean surface pressure have been detected over the past thirty years.

Jaxk Reeves, Detection Of Undocumented Changepoints: Revisions Of The Two-Phase Regression Model

Jaxk Reeves 

Statistics Department, University Of Georgia, USA

and

Robert Lund
XiaoLan Wang

This presentation examines detection and adjustment of climatic series for undocumented changepoint times, primarily from single site data. The two-phase regression model techniques currently used are demonstrated to be biased toward the conclusion of an excessive number of unobserved changepoint times. Simple statistically defensible revisions of these methods based on work by the authors is presented.

G.W. Reuter, Severe convective storms with large hail and F3-F4 tornadoes in Alberta, Canada

G.W. Reuter 

Department Of Earth and Atmospheric Sciences, University of Alberta, Canada

and

M Dupilka

During the past 20 years there have been three tornadic thunderstorms in Alberta with intensities of F3 or greater on the Fujita scale. These events were: the Edmonton tornado of 31 July 1987, the Holden tornado of 29 July 1993, and the Pine Lake tornado of 14 July 2000. A detailed synoptic analysis of these storms was done with the emphasis on surface moisture and storm tracks. All three cases supported the Smith and Yau conceptual model of severe thunderstorm outbreaks in central Alberta. In each case the environment had a capping inversion to build up potential instability, and large values for the convective available potential energy and wind shear. However, the storms differed in the triggering mechanism to break the cap and release the latent instability. The Holden and Pine Lake storms were triggered and developed along a well defined dryline. In contrast, the Edmonton tornado had no dryline; instead the boundary layer was uniformly moist. The three cases also showed differences in the temperature advection. For the Pine Lake storm there was mid-upper level cooling whereas, for the Edmonton and Holden cases there was low level warming but little or no mid-upper level cooling. A comparison of the tracks of the tornadoes showed that the Pine Lake and Holden storms moved consistently in both direction and speed. The Edmonton tornado, however, made an abrupt change in direction and speed. Thus, extrapolation of tornado movement would not be a viable nowcasting technique in this case.

HA Riphagen, Trends in observations for numerical weather prediction

HA Riphagen 

South African Weather Service, South Africa

Numerical weather prediction (NWP) is an initial-boundary value problem in which the governing equations are integrated forward from a fully determined initial state. An optimal, physically consistent initial state, consisting of values of meteorological variables at points of a regular three-dimensional mesh at a particular time, is the product of a data assimilation process in which irregularly spaced observations of many types are blended statistically with short-range forecasts. The more accurate the estimate of the initial conditions, the more accurate the model forecasts.

The NWP systems of the world are currently fed by a wide variety of observing systems, each subjected to stringent testing before being introduced operationally. The results of such testing presented at WMO-sponsored workshops in 1997 and 2000 led to conclusions on the contributions of these systems to large-scale forecast skill at short and medium range. The global observing system has since been significantly enhanced through the launch of several new satellites and instruments. New strategies for observation targeting and the design of optimal observation networks are also enhancing the efficacy of conventional measurements, such as radiosonde and aircraft data. At the same time the emphasis is shifting more and more to the smaller scales as model resolutions increase and 'nowcasting' techniques for severe weather prediction mature. Precipitation is being assimilated at several centres.

Intensive studies have been carried out in global and regional modelling systems to assess the impact of the new data, and of the established data types in the changing observation environment. Further studies have considered the role of well-managed cooperative regional programmes such as EUCOS (Eumetnet Composite Observing System) and AMDAR (Aircraft Meteorological DATA Relay) in facilitating new adaptive observing system concepts designed to shift resources from well-observed to poorly-observed areas. Results and recommendations from the recent Third WMO Workshop on the Impact of Various Observing Systems on Numerical Weather Prediction, where such studies were the focus, will be reviewed with emphasis on the contrasting needs of more and less observation-affluent regions of the world and, in particular, the southern African situation.

JW Roberts, Spatio Temporal Correlation Analysis of NDVI in the Succulent Karoo

JW Roberts 

CSAG, University of Cape Town, South Africa

and

Bruce Hewitson

Emma Archer

Chris Jack

The use of the Normalized Difference Vegetation Index in studies of the relationship between vegetation and rainfall are widespread. Both the spatial and temporal relationships between neighbouring pixels in the succulent Karoo biome were analysed using a Spatial Temporal Correlation Filter (STCF). The STCF calculates the correlation of NDVI values through time with those of the surrounding pixels through time. The mean correlation value is then assigned to the centre pixel. The filter moves onto the adjacent pixel and repeats the process. Two kernel sizes were used 3X3 and 5X5. Results from each are reported. The analyses attempted to identify areas that change through time in a similar way.

The succulent Karoo biome of southern Africa is part of the greater Cape floral Kingdom. Levels of endemism and species richness in the biome are unrivalled. Land use practices namely cropping and grazing have denuded large areas in the biome. Continued agricultural expansion and overstocking pose major threats to the biomes floral diversity. Assessing the anthropogenic impacts and climate related changes imposed on the biome are especially important. Using NDVI and Interpolated rainfall data to assess these impacts provides extremely valuable data.

Time series analysis of NDVI and interpolated rainfall data are usually performed on the premise that rainfall is the dominant driving force in seasonal vegetation change. In the succulent Karoo this premise may not stand. Land use, irrigation of croplands and the influence of river systems receiving their rainfall at times other than the dominant winter rainfall regime all hinder the analysis. It is therefore important to know where seasonal vegetation response is both spatially and temporally homogenous

Analysis was performed for the time period 1985-2003 and for various seasonally defined periods: DJF, MAM, JJA & SON. Results from both the 3X3 and 5X5 Kernel were stratified into classes containing the following correlation values, 0.91 or stronger (class 1), 0.9 – 0.71 (class 2), 0.7 – 0.51 (class 3) and –0.5 – 0.5 (class 4). Class 1 identifies largely undisturbed areas of the lowland plains of the Namaqualand region. Class 2 correspond to areas with increased elevation and rough topography. Class 3 may be classified as a transition zone between class 2 and class 4 as the two classes are found in close proximity to each other. Class 4 is associated with irrigated agriculture and areas around major river systems such as the Orange and Olifants rivers.

Results indicate that large areas of the biome show relatively good correlation values; these are however variable depending on the size of the kernel used and time period in question. Results will contribute to a statistical analysis of the relationship between rainfall and vegetation growth on an intra seasonal time scale

Alan Robock, Detection Of The Effects Of Volcanic Eruptions On Climate

Alan Robock 

Department Of Environmental Sciences, Rutgers University, USA

and

Georgiy Stenchikov

Large tropical volcanic eruptions, such as the 1963 Agung, 1982 El Chichon, and 1991 Pinatubo eruptions, have a detectable influence on global climate. These effects include tropical cooling, but only summer cooling with winter warming in the midlatitudes of Northern Hemisphere. These climate response is modulated by the phase of the quasi-biennial oscillation at the time of the eruption and includes the indirect impacts of volcanically-induced ozone depletion. The effects of smaller eruptions with troposphere-only emissions or sulfur-poor stratospheric emissions, such as the 1980 Mt. St. Helens eruption, are not detectable. High latitude eruptions, such as the 1783 Laki eruption, produce quite different effects. This talk will discuss the ways that volcanic eruptions perturb not only the radiative balance of the planet, but also atmospheric dynamics, and describe the patterns that have been detected with observations and simulated with global climate models. Detection of mid- and high latitude influences, particularly in the winter, requires a large number of ensemble members to overcome weather noise. Our recent simulation of the effects of the Pinatubo eruption, including the effects of the quasi-biennial oscillation, used 24-member ensembles and was able to successfully model the climate response during the next two Northern Hemisphere winters.

M Rouault, INTENSITY AND SPATIAL EXTENSION OF DROUGHTS IN SOUTHERN AFRICA SINCE 1901

M Rouault 

Department Of Oceanography, University of Cape Town, South Africa

Due to the complexity of rainfall regime in Southern Africa, it is difficult to monitor drought in real time with a chart showing percentage from normal or anomaly in total rainfall. Drought occurs often in Southern Africa in all climatic areas at all times of the year with different intensity, spatial extension and duration. Agricultural drought (3 to 6-month time scales) and hydrological droughts (1 to 2-year scale) are often decoupled. The standardized precipitation index (SPI) is a simple index based on rainfall only that can measure drought at different time scales (3, 6, 12 and 24 months) with spatial homogeneity. The SPI is linked to the probability occurrence of dry or wet events. The SPI allows monitoring operationally any location with a 30-year time series. The SPI can also be used to monitor excess moisture during a wet season that can aggravate consequences of floods. This may indicate a saturated catchment conducive to flood development if rainfall continues. It was recently used in many parts of the world and South Africa but never in Southern Africa as a whole. This first application on the 0.5x0.5 degree resolution 1901 1999 rainfall dataset developed by Mitchell et al (2004) at CRU permit to observe some spatial and temporal characteristics of the droughts. Furthermore, their distribution is particular with: The strongest droughts have happened during the mature phase of El Niño

There is considerable decadal variability in the spatial extension and intensity of droughts. Droughts lasting 3 years are not uncommon

Behrooz Sari Sarraf, Evaluation Of Wet And Dry Periods In North West Of IRAN

Behrooz Sari Sarraf 

Geographic Department, Tabriz University, Iran

and

Reza Raihany nia

Abstract: In this paper evaluation of wet and dry periods of southwestern region of Iran (Aras and Urmia basins) is carried out using graphical representation and calculation based on mathematical modeling. Monthly precipitation records of twenty stations for a period of 33 years, from 1970 to 2002 were chosen and fluctuations of each period with respect their numbers were plotted on a semi -logarithmic scale and the regression equations were obtained. Accordingly since the longest wet and dry period happens only once in each period, it is possible to estimate the durations of wet and dry periods for each station using the semi logarithmic graphic representation mentioned above.

Simon Christian Scherrer, Trends In Swiss Alpine Snow Days - The Role Of Local And Large Scale Climate Variability

Simon Christian Scherrer 

Swiss Federal Office Of Meteorology And Climatology (MeteoSwiss), Switzerland

and

Christof Appenzeller

Swiss Alpine snow cover is varying substantially on interannual to decadal time scales. Since the late 1980s a large decrease in snow days has been observed for stations below 800m altitude. In this work multiple linear regression models are used to quantify the importance of seasonal mean local temperature and precipitation to explain the observed variability and trends in Swiss Alpine snow days for the period from 1958 to 1999. The role of large-scale European-North Atlantic climate variability is also discussed. Results show that local as well as large-scale predictor models can account for a modest fraction of ~32 to 45% of the observed snow day variability. However, it appears that at low locations the recent decrease in snow cover can be mainly attributed to an increase in seasonal mean temperature. Seasonal mean precipitation does neither explain large amounts of variability nor affect recent trends in a substantial manner. Differences are found for the northern and southern parts of Switzerland concerning the influence of large scale climate modes. Interannual snow day variability of the northern Swiss Alpine slopes is almost unaffected by the North Atlantic Oscillation (NAO). Decadal trends however can be explained via temperature only by a model that includes the 'NAO index' as explanatory variable. For the southern Swiss Alpine region the NAO is not just explaining the decadal scale trends but also a substantial part of interannual variability via temperature and to a small degree via precipitation. In respect of global warming, these results suggest that a possible change to more frequent positive NAO winters could amplify the decrease in low altitude snow cover much more than expected from an averaged global temperature increase alone.

Jürg Schmidli, Statistical Downscaling Using GCM Precipitation As A Predictor. An Evaluation For The European Alps

Jürg Schmidli

Atmospheric And Climate Science ETH, Switzerland

Conventional statistical downscaling methods for precipitation are prone to problems of 'missing' sensitivities, i.e. sources of variability not represented in the chosen predictors. Using GCM simulated precipitation as a predictor is likely to minimize such problems. This study evaluates the potential of using GCM-simulated precipitation downscaling in the European Alps. The evaluation is based on reanalyses (NCEP, ERA40) which are considered as quasi-perfect GCMs and allow comparison of interannual variations from downscaled results and observations. Four statistical downscaling methods operating on a daily time scale are investigated. Two methods require GCM simulated precipitation as their only predictor. Two more sophisticated methods require in addition some proxy of the large-scale flow. The simple downscaling methods can serve as a benchmark for more complicated downscaling methods such as neural networks or regional climate models. The downscaling methods are evaluated over the European Alps on a seasonal basis for the 15 years 1979-1993. The evaluation statistics encompass several diagnostics, including mean precipitation, and statistics representative of heavy precipitation and drought spells. In addition to biases, we put some emphasis on the representation of observed interannual precipitation variations which are considered as a manifestation of climate sensitivities. The highest skills are obtained for the winter season, when the precipitation climate is driven by synoptic-scale disturbances. The lowest skills are found for the summer season, when local convection is more dominant. The results of this study imply that GCM-simulated precipitation can carry interesting information on regional precipitation variability even in a region of complex terrain.

David Sexton, Constraining Probabilistic Climate Predictions By Model Reliability

David Sexton 

Hadley Centre For Climate Prediction And Research, Met Office, United Kingdom

and

Mat Collins

Glen Harris

James Murphy

Mark Webb

Uncertainty in climate prediction is not only due to uncertainty in future emissions, atmospheric composition and natural variability but also due to the way climate models represent the real world. To explore the modelling uncertainty, we have produced a 'physics ensemble' of 53 atmosphere models coupled to a thermodynamic ocean 'slab' where each member differed from the Hadley Centre's standard slab model, HadSM3, by a perturbation to one of 29 physics parameters. These slab models, which are relatively quick to run to equilibrium, are forced with 1x and 2x pre-industrial CO₂ levels. From these runs, we estimated the equilibrium annual mean global mean temperature response to doubling CO₂ levels, often called climate sensitivity. For this model and experimental set-up, the relative probability for different values of climate sensitivity can be estimated. A problem with this calculation is that the ensemble members differ in quality and reliability; IPCC TAR (The Intergovernmental Panel on Climate Changes Third Assessment report) faced a similar problem basing their uncertainty range on future warming on models from different climate centres. We therefore develop a Climate Prediction Index, based on the root mean square errors of 32 model 20-year means from suitable observational mean data sets, to measure the relative quality of the ensemble members. Weighting by model reliability changes the 90% confidence interval from 1.8-5.7K to 2.4-5.7K. By doing this, we hope to give more weight to climate models that best represent the important physical feedbacks for climate change. As the mean climate is not necessarily a direct measure of how well a model represents these feedback processes, we show some preliminary results investigating the extent to which the mean climate constrains future climate change.

Leonard Smith, PROBABILISTIC SEASONAL FORECAST IN THEORY AND PRACTICE

Leonard Smith 

Centre for the Analysis of Time Series Department of Statistics, London School of Economics,
United Kingdom

The DEMETER project has successfully produced multi-model multi-initial condition seasonal forecasts with a lead time of six months over a period of several decades. After an overview of the available hindcast dataset, consisting of nine initial conditions each evolved under seven global coupled atmosphere-ocean models initialised four times each year, a number of questions on the interpretation, evaluation and application of the DEMETER dataset are addressed. It is argued that these three issues are fundamentally linked; issues as basic as the "correct" approach to "bias correction" will have fundamental implications not only on forecast evaluation and ensemble design, but also on the utility of the ensemble forecast system in application. Central to these questions is the clear distinction between two goals: does one view the ultimate goal of a multi-model multi-initial condition ensemble prediction system as providing insight into some hypothetical "True" probability distribution function of likely futures, given our current limited knowledge? or is the goal to extract the probability distributions of each of the available models and their relation to the single realization target (which might be "Truth", but in the case of DEMETER is an ERA-40 reanalysis). It is argued that, ideally, one would condition a seasonal forecast on the joint distribution of all forecast information (admitting that this argues for the inclusion of models with relatively 'low skill' but 'high information' content). Practical applications, however, require practicable methods; current deployable methods in the agricultural and health sectors which were used within DEMETER are noted, and the search for consistency tests to evaluate proposed methods is reported.

Leonard Smith, Ensemble Climate Modeling: WHAT Can IT Teach Us Within An in-sample Science

Leonard Smith 

Centre for the Analysis of Time Series Department of Statistics, London School of Economics,
United Kingdom

By the nature of the problem addressed, climate modelling is an in-sample science. This is true on at least two counts: first, the time-scale of the forecasts is significantly longer than the lifetime of the simulation model(s) used; second, both the transient nature of the boundary conditions (external forcing and atmospheric composition) and slow rate at which new data is obtained limits the scope of traditional approaches like cross-validation. There are fundamental limitations to what we can be sure of, even on what we can assign meaningful probabilities to, in this situation (see L.A. Smith, (2002) Proc. National Acad. Sci. USA, 4 (99): 2487-2492). If we are to realise the potential of ensemble experiments, it is important to clarify the questions the experiment is to address at the design stage. Several competing goals are discussed in the context of multi-model, multi-parameter-value, multi-initial condition ensemble modeling. It is critical to distinguish (at the design stage) between (a) making the best forecast possible today, in the knowledge that it may be shown to be physically irrelevant (in, say, 10 years time) well before the verification time; (b) investigating the behavior of today's model(s) so as to provide a more useful forecast model(s); (c) bounding likely future behaviours (that is, giving probable limits under each model, without a clear indication of their probability in practice) and (d) suggesting specific, pivotal observations to allow repeated reinterpretation the forecast range (using past and suggesting future model simulations) as time passes. The common disparity between the variables of interest in potential application (regional, seasonal, and extreme) and the variables most discussed in model evaluation (global, annual averages) is used to illustrate the importance of a true initial condition component within the ensemble mix for climate studies in each case above.

Tim Staeger, Statistical Separation Of Natural And Anthropogenic Signals In Observed Surface Air Temperature Time Series

Tim Staeger 

Institute For Meteorology And Geophysics, J.W. Goethe University, Germany

and

Christian-D. Schonwiese

In this study observed surface air temperature time series are statistically analysed with respect to natural and anthropogenic forcing. The data represent global or hemispheric averages, respectively, as well as regional data (areas means on a global scale, European grid-point data). The period considered covers approximately the recent century. The analysis method applied is stepwise regression, including EOF techniques and moving analysis in time, where low-dimension regression models are evaluated explaining a maximum of observed variance. Thereby, natural forcing is represented by variations of the solar constant, explosive volcanism, El Nino (ENSO), and the North Atlantic Oscillation (NAO), anthropogenic forcing by CO₂ equivalents (greenhouse gases, GHG) and tropospheric sulphate (SU). On a global or hemispheric scale these forcing parameter time series explain c. 65 – 80 % of the observed temperature variance where greenhouse gases dominate followed by solar activity forcing. The SU signal appears to be problematical. The way to a more regional consideration (as well as the analysis of precipitation and air pressure data) leads to an increase of stochastic variability (noise) and in turn to both less explained variance and less confidence of the GHG signal. Although the most pronounced GHG signals are found in the mid-continental areas of the northern hemisphere, maximum confidence is related to oceanic and tropical regions. It is concluded that the large-scale average, in particular global, of surface air temperature is most appropriate to detect anthropogenic forcing on climate by means of observations.

David B. Stephenson, An Overview of EOF / PCA Techniques in Climate Research

David B. Stephenson 

Department of Meteorology, University Of Reading, United Kingdom

This talk will provide an overview of Empirical Orthogonal Function (EOF) techniques used in climate research. Since the 1950s, EOF analysis (or Principal Component Analysis) techniques have been widely used in climate analysis. This talk will attempt to address the following eight questions:

- What is EOF/PCA analysis?
- Why do climate scientists use these techniques?
- How has EOF/PCA research evolved since 1950?
- What exactly is a physical/dynamical mode?
- What is a modal expansion?
- Which is more physical: the North Atlantic Oscillation or the Arctic Oscillation?
- What about more regional modes in the Northern Hemisphere?
- Where are we going next?

It is hoped that this talk will raise several issues for further discussion and in so doing will capture much of the variance in current EOF thinking!

Dáithí Stone, The End-To-End Attribution Problem: From Emissions To Impacts

Dáithí Stone 

AOPP, Department Of Physics, University Of Oxford, United Kingdom

and

Myles Allen

The attribution of recent changes in global mean climate to anthropogenic activities is now well established. Of course, it is not global mean climate but rather local episodic extreme weather events that often have the largest impacts. While the attribution of climate change at more regional scales is now being examined, this research is still restricted to continuous quantities. Here we present a transfer of a methodology used in epidemiology for the attribution of episodic events. Key to this approach is the consideration of the risk of the event occurring, rather than the occurrence itself. Given restrictions on our knowledge of the climate system, it is important to recognise that this risk is in fact a probabilistic quantity and should be treated accordingly. Tests of the application of this technique to the simple Lorenz 1963 system will be examined.

Dáithí Stone, What Is The Best Current Predictor Of Future Climate?

Dáithí Stone

AOPP, Department Of Physics, University Of Oxford, United Kingdom

and

Jesse Kenyon

Myles Allen

Gabriele Hegerl

Dave Stainforth

Predictions of climate change produced by general circulation models can only be verified against historical observations. However, due to sampling issues the observational constraints are consistent with a wide range of possible futures. Thus, for establishing the robustness of the prediction of some future variable we must examine the full historical climate variable space. It is of considerable interest therefore to establish which direction of that variable space is most important, both in order to constrain the future prediction and to focus efforts in model development. Here we conduct a modelling study using the CMIP2 quasi-ensemble to investigate how intermodel variations in future predictions relate to variations in the reproduction of historical quantities. Details of the technique and the modelling tests will be presented.

Peter Stott, Detection Of Anthropogenic Climate Change In Africa And Other Continental Regions And Attribution Of The Increasing Likelihood Of Extremely Hot Summer Temperatures.

Peter Stott 

Hadley Centre For Climate Prediction And Research, Met Office, United Kingdom

and

Daithi Stone
Myles Allen

We show that anthropogenic increases in annual mean temperatures have been detected on individual continents including in Africa, North America, and Europe. Motivated by the extremely hot summer in Europe in 2003, we examine the increasing likelihood of hot European summers and show that a large fraction of the risk of very hot summers can already be attributed to anthropogenic causes.

**Matthew Swann, Extreme Winter Wind And Precipitation In North-Western Europe:
Dataset Comparison And Interannual Variability.**

Matthew Swann 

Climatic Research Unit, University Of East Anglia, United Kingdom

and

Tom Holt

This work describes the development stages of a seasonal forecasting model designed to predict daily extremes of wind speed and rainfall over north-western Europe in winter. European extremes of wind speed and rainfall are embedded in the mean westerly extratropical circulation and are, therefore, influenced by a number of large-scale processes which are persistent on seasonal timescales. These potential predictors include, for example, the North Atlantic Oscillation (NAO), the sea surface temperature field over the North Atlantic, and upper air circulations (e.g. Thompson et al., 2002; Frankignoul et al., 2003). The predictor processes are partially coupled, and have been shown to vary on interannual timescales (e.g. Rodwell and Folland, 2002). Before considering predictor/predictand relationships, it is essential to perform a detailed study of the predictands. This preliminary analysis determines the optimum statistical model, helps identify the most appropriate methodology, and is the basis of this discussion. The predictands are total daily precipitation and maximum daily wind speed, taken from station records and the ECMWF ERA40 reanalysis. The station data are used to assess the extent and nature of smoothing of extremes in the reanalysis data. First, the statistical properties of each variable from the two datasets are compared by creating frequency distributions at selected coincident points. To assess extremes, the analysis compares the properties of the Generalised Extreme Value distribution for the variables in each dataset at the chosen points. The overall interannual variability of the datasets is examined by comparing Principal Component Analysis (PCA) scores for leading modes of the precipitation and wind fields.

Mark Tadross, Uncertainty In Producing Regional Climate Change Scenarios For Southern Africa

Mark Tadross 

CSAG, University of Cape Town, South Africa

Using a regional model to downscale climate change predictions from GCMs involves several aspects of uncertainty: GCMs have different biases when simulating the climate, the correct physical schemes to use with an RCM are unclear, as is the correct representation of the land surface in an RCM. All the above are further complicated by a severe lack of observational data with which to test these models. The talk will address these issues in an effort to deal with some of these uncertainties.

Mark Tadross, Midlatitude Wave Patterns Associated With The Start Of The Growing Season In Southern Africa.

Mark Tadross 

CSAG, University of Cape Town, South Africa

The onset of the growing season in southern Africa is an important seasonal characteristic. Evidence is presented for the influence of synoptic variability on onset. Mid-tropospheric height anomalies are classified using self organising maps and positive anomalies to the south and east are shown to be associated with an early start. GCM simulations of these anomalies are also compared to the observations and the implications for forecasting discussed.

Warren James Tennant, The Importance Of Rainfall Characteristics In Forecasting And Verification

Warren James Tennant 
South African Weather Service, South Africa

Spatial and temporal variability of rainfall is difficult to quantify. Rainfall observation techniques all suffer from inadequacies in one way or another such that it is difficult to establish the spatial integral of rainfall in a specific area. Gauges only provide spot measurements, while satellite and radar measurements are difficult to calibrate for different synoptic situations. This study looks at a large number of rain gauges in the Pretoria area to investigate rainfall characteristics on daily and monthly time-scales. The aim is to determine to what extent in-homogeneity in daily convective-type rainfall is transferred to the longer time-scales and how these high-density measurements match up with satellite and radar measurements. The effect of rainfall observations on forecast model development and the verification of rainfall forecasts is also investigated. The period of study for this case is January to March 2004.

Warren James Tennant, Downscaling GCM Forecasts To Intra-Seasonal Rainfall Characteristics

Warren James Tennant 
South African Weather Service, South Africa

Seasonal forecasts typically only provide outlooks of rainfall and temperature anomalies averaged over the entire season. However, there is increasing pressure from users of seasonal forecast guidance for intra-seasonal rainfall characteristics such as rain days, length of dry spells etc. Forecast frequencies of daily circulation archetypes, determined using a SOM analysis, are related to rainfall characteristics using neural networks. This methodology and some results will be presented.

Anneke Thackrah, Climate Change: Is The Brown Locust Trying To Tell Us Something?

Anneke Thackrah 

ARC - Institute For Soil, Climate And Water, South Africa

and

Francois Engelbrecht

Hannes Rautenbach

Margaret Kieser

The brown locust, *Locustana pardalina* (Walker) is endemic to the semi-arid Karoo regions of South Africa and southern Namibia, and is a constant threat to food security within the southern African region. As part of a project to develop a Brown Locust Early Warning System (BLEWS), the locust breeding region was defined using brown locust control data for the period 1984 to 2001. A comparison of the brown locust breeding regions defined in 1937 (Faure), 1958 (Lea) and 2001 (Kieser) reveals a significant westward shift in the eastern boundary. During the period 1984 to date, limited hatching was reported in the eastern Karoo, which influenced the subsequent closure of the Middleburg poison depot and its relocation to Upington. In contrast, the southwestern limits of the outbreak region appear to have expanded. Simulations of climate change by two high-resolution regional climate models (DARLAM and C-CAM), forced by fully coupled Ocean-Atmosphere General Circulation models (AOGCMs), provide much insight into the observed changes in outbreak regions. Modelling results indicate that a general westward shift of cloud bands that normally produce summer rainfall over the western and central interior of Southern Africa can be expected in response to Global Warming. A general increase in annual rainfall across the Karoo can indeed be observed at various weather stations from the early 1900s, with an associated westward shift of the winter and early summer (July - December) 150 mm isohyete. This corresponds almost exactly to the westward shift in the eastern boundary of the locust outbreak area, and suggests that the wetter conditions are unsuitable for brown locust breeding (increased vegetation cover reduces the availability of suitable oviposition sites). In contrast, the expansion of the brown locust outbreak area in the drier southwest (100-150 mm) may reflect enhanced breeding success due to improved soil moisture conditions for egg hatching.

Amanda Townsend, Tracking The Evolution Of UK Extreme Temperature Trends

Amanda Townsend 
University Of East Anglia, United Kingdom

and

Jean Palutikof
Gareth Janacek

As part of a project to understand the changing character of extreme temperature events over the British Isles during the twentieth century, long time series of observed daily maximum and minimum temperatures have been collected. The selection of records has been made bearing in mind the need for good geographical spread. Each time series has been tested for inhomogenities and these have been removed where necessary. The occurrence of extremes has been explored by fitting Generalized Extreme Value (GEV) distributions to annual maxima and minima drawn from the winter (DJF) and summer (JJA) seasons. Thus, the time series of extremes for analysis are: highest and lowest values of winter daily maximum temperatures; highest and lowest values of winter daily minimum temperatures; highest and lowest values of summer daily maximum temperatures; highest and lowest annual values of summer daily minimum temperatures. In each case, one value is drawn from each three month time period. The technique of Maximum Likelihood Estimates has been used to estimate the GEV parameters. A 30-year moving window has been applied to each of the eight time series, progressing by one year at each time step. The parameters of the GEV distribution fitted to each 30-year segment have been calculated and used to determine 20-year return values. The 30 year period ensures a stable and sufficiently large enough dataset in order that unbiased parameter estimates can be calculated. The moving window approach allows exploration of changes though time, smoothed for clarity. From the GEV parameters, 20-year return values have been calculated for each station and then analyzed for trends. While none of the resulting series are precisely linear against time, linear regression offers a useful foundation for trend detection. The magnitude and direction of the trends have been determined using linear regression with the statistical significance determined from the Kendall-Tau test. With the exception of winter extreme cold minimum temperatures (which show a cooling trend of between $-0.83^{\circ}\text{C}/\text{century}$ and $-3.22^{\circ}\text{C}/\text{century}$) all of the time series display some evidence of a warming trend. The six longest station records for cold tail summer minimum temperatures display the most consistent rise in extreme temperatures during the last century with trend magnitudes of $1.59^{\circ}\text{C}/\text{century}$ to $3.18^{\circ}\text{C}/\text{century}$. Understanding the possible causes of these trends will form the next stage of the study. As a first step, certain covariates are to be added into the GEV analysis to establish their likely effects on these trends, these include sea level pressure and specific humidity. Results from this analysis will be presented at the meeting.

Silke Trömel, A Model Fitting Analysis Of Monthly Precipitation Data

Silke Trömel 

J.W. Goethe University, Department Of Meteorology and Geophysics, Germany

and

Christian-D. Schoenwiese

A method for statistical modelling of climate time series and its application to monthly precipitation totals of the 20th century is presented. The statistical model describes the time series under consideration by non-stationary probability density functions with two time-dependent parameters. It depends on the precipitation regime, whether the Gumbel or Weibull distribution achieves complete description of the data. Thereby the location and scale parameter or the scale and shape parameter, respectively, may vary containing structured components like linear or non-linear trends, constant or significantly changing annual cycles and episodic components. So reliable estimates of these structured components can be evaluated and the analytical description of the time series allows probability assessments of extreme values for every time step of the observation period as well. In this way observational German precipitation data can be explained as a realization of a Gumbel-distributed random variable. Mainly in western Germany positive trends in both the parameters of the distributions are detected. Together with changes in the amplitude and the phase angle of the annual cycle in the scale parameter they reflect seasonal differences in the probability development of exceeding, for example, the 180 mm monthly total precipitation threshold. Winter and spring months show positive changes while the late summer months reveal negative changes or a slight increase of this probability.

R.P Tshikalanke, Spatial and Temporal Variations in Domestic Biofuel Consumption rates in Southern Africa.

R.P Tshikalanke 

University of Witwatersrand, South Africa

Biomass burning leads to the production of trace gases and aerosols, which influence the atmospheric chemistry. The magnitude of these emissions is comparable to those of fossil fuel. Studies have indicated that four types of biofuels are used in Southern Africa, namely wood, agricultural residues, charcoal and animal dung. The amount of biofuel consumed at a single part in time has been measured at representative sites around Southern Africa. The seasonal variation of biofuel consumption has, however not been quantified. Three households were selected for long term data monitoring within Southern Africa. The occupants were given weighing scales and logbooks were they are recording the number of kilograms used everyday for three cooking session (breakfast, lunch and dinner). The study aims to investigate the spatial and temporal variation in biofuel consumption rates throughout Southern Africa. It will also investigate environmental factors influencing the use of domestic fuel. This data, together with population data will be used to calculate biofuel consumption rates for Southern Africa.

Hans Von Storch, The Philosophy of statistics

Hans Von Storch

GKSS Research Centre, Institute For Coastal Research, Germany

No Abstract Information Available

Hans Von Storch, Limits Of Historical Reconstructions Using Indirect Evidence

Hans Von Storch

GKSS Research Centre, Institute For Coastal Research, Germany

We test how much variability at centennial timescales is lost when past climates are reconstructed from proxy data. We use data from an atmosphere-ocean climate model simulation of the last 1000 years, forced with estimates of natural and anthropogenic forcings, and the method of Mann et al. (1998) (1) to estimate Northern Hemisphere temperature from grid-point temperatures (with white noise added) in realistic locations. We find that the low frequency variability is greatly underestimated when the predictors contain 50% or more noise. The inclusion of more predictors from Africa and Asia results in only minor improvement.

Jin-Song Von Storch, On Statistical Dissipation In GCM-Climate.

Jin-Song Von Storch

Meteorological Institute / University Of Hamburg, Germany

This work deals with the dissipation that affects the statistical behavior of a climate variable. The hypothesis is tested that the statistical dissipation of large-scale variables in a climate model is significantly influenced by the temporal variations of the model's small scale variables. The test is done with the T21 ECHAM4 atmospheric GCM by varying model's representation of small-scale variables in two different ways. First, the strength of the horizontal diffusion is modified. Secondly, white noise unrelated to the state of the large-scale variables is added to variables with the smallest scales (wave numbers equal to and smaller than 18). It is found that the statistical dissipations of large-scale variables (e.g., vorticity at wavenumber equal to and larger than 6) depend on the intensity of small-scale fluctuations, no matter whether they are induced by modifying the horizontal diffusion or by adding noise. The stronger the small-scale fluctuations, the stronger are the dissipations of the large-scale variables. This result suggests that the simulation of low-frequency climate variations and the prediction of climate change responses depend on the model representation of small-scale climate components.

NM Walton, Characterization of the Cape Town Brown Haze

NM Walton 

University Of Witwatersrand, South Africa

and

S Piketh

Visibility impairment occurs as a result of the scattering and absorption of light by particles and gases in the atmosphere. It is described as the haze which obscures the clarity, colour, texture, and form of what is seen through the atmosphere. Primary particles, secondary particles (sulfates and nitrates), water particles and nitrogen dioxide gas can cause significant atmospheric visual impacts (Latimer and Samuelsen, 1978). Cape Town, a major local and international tourist attraction, experiences high pollution levels during winter with the occurrence of a brown smog or 'brown haze'. To evaluate the impact of the haze on visibility impairment, the Cape Town Brown Haze II project was undertaken during July and August 2003. The plume visual impact screening model, VISCREEN, was used for an initial data analysis. Two screening criteria were used to assess the potential impacts: plume contrast and ΔE . Plume contrast is the relative difference in light intensity of two view objects while ΔE is a measure of the perceived difference in brightness and colour between a plume and the background over a range of colours in the visible light spectrum (USEPA, 1992). The plume visibility model, PLUVUE II, was then used to model plumes from the Table View and Bellville areas to calculate visual range reduction and atmospheric discolouration over the surrounding areas.

Latimer, D.A and Samuelsen, G.S., 1978: Visual impact of plumes from power plants: A theoretical model, Atmospheric Environment, 12, 1455-1465.

U.S. Environmental Protection Agency., 1992: Workbook for Plume Visual Impact Screening and Analysis (revised), EPA-454/4-88-015, October 1992, U.S. Environmental Protection Agency, Research Triangle Park, NC.

Xiaolan Wang, Using Non-Stationary Generalized Extreme Value Models To Assess Historical And Possible Future Changes Of Extreme Wave Heights In Northern Hemisphere Oceans

Xiaolan Wang 

Climate Research Branch, Meteorological Service Of Canada, Canada

and

Mr. Val Swail

In this study, we review and extend previous studies on changes of wave heights, focusing on both observed and possible future changes in the North Atlantic and in the North Pacific. Two wave hindcasts are used here as observed waves for 1958-1997: a global wave hindcast based on the 10-m winds taken from the NRA (i.e., the reanalysis of the National Centers for Environment Prediction, National Center for Atmospheric Research), and a detailed North Atlantic wave hindcast produced with intensively reanalyzed surface winds over the North Atlantic basin. The observed relationships between sea level pressure (SLP) and significant wave height (SWH) are used to construct climate change scenarios of SWH: Projections of seasonal extreme SWH are made using non-stationary generalized extreme value (GEV) models. The GEV models are trained using the NRA SLP and the observed SWH data for 1958-1997. The SWH scenarios are constructed using seasonal mean SLP and squared SLP gradient fields from a coupled climate model (CGCM2) under three different forcing scenarios. The projected (and observed) trends/changes in SWH are assessed by conducting a trend analysis, in which linear trends are evaluated against quadratic trends. Both oceans had significant changes in both winter and fall seasonal extremes (and means) of SWH during 1958-1997; they are also projected to have significant changes in the 21st century under all three forcing scenarios. The rate and sign of the projected future SWH changes are not constant throughout the 21st century; and in some regions, these appear to be quite dependant on the forcing conditions. Often, the projected SWH changes are characterized either by faster increases in the late decades than in the early decades, or by decreases in the early decades followed by increases, depending on the forcing scenario and the specific location. The rate of SWH change appears to have a positive relationship with the rate of increase in the greenhouse gases forcing. Changes in ocean wave heights are associated with changes in storm tracks and cyclone activities over the oceans. In the projected warmer climate, the North Pacific storm track rotates clock-wise in winter (JFM), with more frequent occurrence of strong cyclones in the southeast and northwest of the North Pacific; while the storm track over the northeast Atlantic slightly rotates clock-wise in fall (OND), with more frequent occurrence of strong cyclones in northern Europe and over the Norwegian Sea.

Xiaolan Wang, Assessment Of Observational Biases And Trends In Canadian Cloudiness

Xiaolan Wang 

Climate Research Branch, Meteorological Service Of Canada, Canada

and

Dr. Francis Zwiers

Mr. Val Swail

Mr. Ted Yuzyk

We have analyzed the hourly cloudiness record (in tenths of the sky dome) at 103 Canadian stations where observations were made by human observers. The records at 56 stations span the period from January 1953 to April 2003, while the remaining 47 stations have shorter records (35-49 years). All records are nearly complete. These records were first subjected to an intensive assessment of observational biases, and were subsequently analyzed for trend by means of logistic regression. This assessment involved statistical change-point analyses, station metadata investigation and visual inspection of the time series in question. Changes in observing practices and observer training, frequent observer changes, relocation, etc. were found to introduce significant step-changes in cloudiness time series, especially in the period before the latest version of Canadian Manual of Surface Weather Observations (MANOBS) was published in January 1977. We carried out the analyses for each of the 11 cloudiness conditions (from clear sky to overcast: 0/10 - 10/10) separately, while preserving consistency among the 11 categories. The results of our analyses show that observational biases in Canadian cloudiness records are too numerous and too significant to ignore. Some biases are such that trends estimated from unadjusted cloudiness data can be far from reality. After removing the effect of observational biases, the overall trends of Canadian cloudiness features more frequent occurrence of clear sky, nearly overcast (9/10) and overcast conditions, and less frequent occurrence of broken sky conditions in daytime; while nighttime trends feature more frequent medium-heavy cloudiness conditions and less frequent clear sky and light (1-2 tenths) cloudiness. Overall nighttime cloudiness has increased during the past half century, which is consistent with the strong increase in daily minimum temperature that has been observed in Canada.

A.M. Webb, The nature of the upwelling cell south of Madagascar

A.M. Webb

Department Of Oceanography, University of Cape Town, South Africa

and

R. E Lutjeharms

E Machu

Upwelling in the ocean plays a crucial role in the ecology of many coastlines of the world and dramatically affects the weather and climate of the adjacent land masses. Most extensive upwelling regimes are found along the west coasts of continents and are fairly well understood. It is unusual for new upwelling systems still to be discovered. One such newly found upwelling cell is south-east of Madagascar. This upwelling cell was predicted from basic theory and as a result subsequently identified in thermal infrared satellite imagery as well as satellite images of ocean colour (Lutjeharms and Machu, 2000). This has led to a research cruise making specific observations in the region as part of the Dutch/South African ACSEX II expedition on the Dutch research ship Pelagia. The initial results of this cruise confirmed that there indeed is an upwelling process that occurs at this location (Machu et al., 2002). Further analyses have now been undertaken and show the nature of the upwelling as well as the likelihood of the retroflexion of the East Madagascar Current south of Madagascar. It is shown that the East Madagascar Current carries a core of Red Sea Water as it passes the continental shelf off south-eastern Madagascar. Inshore of the current there are indications of cooler and saltier water as a result of upwelling. Satellite and other products suggest that the current generates a cyclone that traps shelf water and carries it away, while an anticyclone forms as part of the retroflexion. This dipole pair then moves off towards the South African coastline (De Ruijter et al., 2004).

De Ruijter, W.P.M., H.M. van Aken, E.J. Beier, J.R.E. Lutjeharms, R.P. Matano and M.W. Schouten, (2004). Eddies and dipoles around South Madagascar : formation, pathways and large-scale impact. Deep-Sea Research I, in press.

Lutjeharms, J.R.E. and E. Machu, 2000. An upwelling cell inshore of the East Madagascar Current. Deep-Sea Research I, 47(12): 2405-2411.

Machu, E., J. R. E. Lutjeharms, A. M. Webb and H. van Aken, 2002. First hydrographic evidence of the southeast Madagascar upwelling cell. Geophysical Research Letters, 29(21), 2009, doi:10.1029/2002GL015381.

R Wilby, Limits To Downscaling Predictability / Skill

R Wilby 

Climate Change Science, Environment Agency, United Kingdom

This talk will draw on a decade of downscaling intercomparison studies to review issues of predictability and skill in downscaling techniques; raising notably the debate of to what degree it is important to select one approach over another. In addition, the talk will present recent developments and results arising from the Statistical and Regional dynamical Downscaling of Extremes for European regions (STARDEX) project which has focused on rigorous and systematic inter-comparison and evaluation of statistical, dynamical and statistical-dynamical downscaling methods.

Xuebin Zhang, On The Estimation Of Exceedance Over A Threshold

Xuebin Zhang 

Climate Research Branch, Meteorological Service Of Canada, Canada

and

Gabriele Hegerl
Francis Zwiers

Indices of climate extremes have been developed and maintained for the purpose of monitoring climate change. They are also potentially useful for climate change detection studies. A number of such indices are defined as the number of days in a year or season that daily values exceeded a time dependent threshold. These thresholds are typically defined as an annual cycle of daily percentiles that are estimated from a base period such as 1961-1990. Time series of the frequency of exceedance above the estimated thresholds are often analyzed for trends. Using Monte Carlo simulation, we demonstrate that these time series may contain artificial jumps at the beginning and end of the base period. This would make these exceedance frequency time series unsuitable for monitoring and detecting climate change. A bootstrap resampling procedure is proposed to estimate exceedance frequencies during the base period effectively removes the inhomogeneity.

Xiaogu Zheng, Interannual Variability Of Seasonal-Mean Fields

Xiaogu Zheng 
NIWA, New Zealand

and

Carsten Frederiksen

We propose a method for studying the influence of intraseasonal variability on the interannual variability of seasonal mean fields. The method, using monthly mean data, provides estimates of the interannual variance and covariance, in the seasonal mean field, associated with intraseasonal variability. These estimates can be used to derive patterns of interannual variability associated with physical processes that vary significantly within a season, such as atmospheric blocking, or intraseasonal oscillations. Removing this intraseasonal component from the total interannual variance/covariance, one can define a “slow” component of interannual variability that is closely related to very slowly varying (interannual/supra-annual) external forcings and internal dynamics. Together these patterns may help in our understanding of the source of climate predictive skill, and also the influence of intraseasonal variability on interannual variability. As an example, the methodology is applied to diagnose physical processes that play major roles in the variability of DJF New Zealand temperatures.

Modes of variation – some introductory remarks

Ian Jolliffe
Universities of Reading, Aberdeen,
Southampton
i.t.jolliffe@reading.ac.uk

What are ‘modes of variation’

- Should they be
 - Defined by the physics
 - Explainable by the physics after the fact
- Or can they be
 - Purely empirical - defined by the mathematical or statistical techniques that find them

How do we find ‘modes of variation’

- Using physics alone (not at this meeting)
- Using descriptive statistical techniques
 - If so, which techniques?
 - The choice depends on what we mean by ‘variation’
 - If variation = variance, then PCA/ EOF analysis is the obvious choice
- Using a model-based statistical approach – a different sort of ‘mode of variation’?

EOFs or not?

- The great advantage of PCA/EOFs is that statistically we know exactly what we are finding even if we sometimes have trouble physically interpreting what comes out
- Drawbacks/restrictions
 - Linearity
 - Physical interpretation (sometimes)
 - Results partly determined by properties of the technique rather than the data (but also true of all(?) competitors)

Competitors to EOFs

- Mainly achieve one of the following
 - Non-linearity
 - Simplification of numerical/physical interpretation (sometimes more numerical than physical)
 - Modes of variation that link two or more data sets
- A key question for any competitor is ‘Do we understand what the technique is doing beyond producing pretty pictures?’

A FEW REMARKS ON PCA TECHNIQUES IN CLIMATE RESEARCH

OTHER APPLICATIONS OF PCA

- e.g., PCA as a classification tool (alternative to cluster analysis)
- “non-standard” applications
 - With fixed time (PCA of trend magnitudes, Huth this Tuesday)
 - With fixed space (relationships between climate elements at a single station; prefilter prior to weather type / air mass classification)

SOME METHODOLOGICAL REMARKS

- Careful selection of methodological options
 - mode (2 choices for fixed variable; S x T)
 - S-mode doesn't produce “circulation types”!!!
 - similarity matrix
 - rotation

SOME METHODOLOGICAL REMARKS

- Interpretation of results to be done \pm careful check: is output real or artifact?
 - Several ways to check it, including:
 - Stability of rotated PCs
 - Simple structure
 - Composite analysis
 - Congruence PC loading x corresponding correlation map

PCA in detection of atmospheric circulation modes

- Non-linearity ! (Metelka later today)
- Verification of reality: mode compositing
 - NAO x AO: Cash et al. (JAS 2002) – modes on aquaplanet
 - PC1 annular
 - But: composites of PC1 high-amplitude events two-sausage-like (= NAO-like)
 - \pm “AO” doesn't occur in reality
 - Shorter averaging period than 1 months


A regional Intensity-Duration-Frequency curves based on precipitation scaling properties in Belgium

D. Gellens

9 IMSC



Introduction and purpose



Presentation outline


- § Introduction and purpose
- § Observation networks
- § Extreme values distribution
- § Regional approach for estimating at-site fractiles
- § Modeling the I.D.F. curves
 - 1 Modeling the regional growth curves parameters
 - 1 Scaling properties of extreme means
 - 1 Modeling the extreme means
- § Practical results
- § Conclusions

9 IMSC



Observation networks

- § Daily precipitation (8h-8h)
- § Long-term records
- § Aggregation time steps: 1, 2, 3, 4, 5, 7, 10, 15, 20, 25 and 30 days
- § 10 min recording rain gauges
- § Aggregation time steps: 10 min, 20 min, 30 min, 1 h, 2 h, 6 h, 12 h, 24 h, 48 h, 72 h, 96 h, 120 h, 144 h and 168 h.



Extreme values distribution

- § GEV distribution

$$F(x) = \exp \{ - (1 - \theta(x - \xi) / \alpha)^{1/\theta} \} \text{ with } \theta(x - \xi) / \alpha < 1,$$

ξ is the location parameter, α the scale and θ the shape


- § for $\theta=0$ GEV corresponds to a Gumbel distribution
- $$F(x) = \exp \{ - \exp (-(x - \xi) / \alpha) \}$$

9 IMSC



Regional approach for estimating the at-site fractiles

- § Regional homogeneity of the L moments ratios (Hosking and Wallis, 1997)
- § Adequation test according to various distributions e.g. GEV, lognormal, Pearson type III, ... (Z^{DIST} tests)



Regional GEV distribution parameters

$$Q_{kTs} = \xi_{ks} + \frac{\alpha_{ks}}{\theta_{ks}} \{1 - (-\ln(1-1/T))^{\alpha_{ks}}\}$$

$$Q_{kTs} = \mu_{ks} \cdot x_{kTr}^R = \mu_{ks} \cdot \left(\xi_k^R + \frac{\alpha_k^R}{\theta_k^R} \{1 - (-\ln(1-1/T))^{\alpha_k^R}\} \right)$$

- § PWM estimators (L-moments)
- § at-site means μ_{ks} , regional L-CV τ^R and regional L-skewness τ_3^R (Hosking et al., 1985)
- § $\theta_k^R = \theta_k^R(\tau_3^R)$
- § $\alpha_k^R = \alpha_k^R(\theta_k^R, \tau^R)$
- § $\xi_k^R = \xi_k^R(\alpha_k^R, \theta_k^R)$ as the mean = 1

9 IMSC



Regional GEV distribution parameters


$$\text{var}\{Q_{kTs}\} = \mu_{ks}^2 \cdot \text{var}\{x_{kTr}^R\} + \text{var}\{\mu_{ks}\} \cdot (E\{x_{kTr}^R\})^2$$

- § Asymptotic expressions of the variance of the PWM fractile estimates (Gellens, 2002) while combining the regional approach with data extension procedure
- § Comparison with other procedures


9 IMSC



Regional growth curves



At-site fractiles



Modeling the regional growth curves parameters

- § $\theta = \theta(k)$ and $\alpha = \alpha(k)$


$$\bar{x}_{kTr}^R = 1 + \frac{\bar{\alpha}^R(k)}{\bar{\theta}^R(k)} \left[\Gamma(1 + (\bar{\theta}^R(k))^k) - (-\ln(1-1/T))^{\bar{\theta}^R(k)} \right]$$

- § Purpose:
 - 1 express the fractiles as a function of k and extend the range of k values
 - 1 combine information from the two networks


9 IMSC




Parameter α



Parameter θ



Scaling properties of extreme means




9 IMSC



Modeling the at-site means

- § Fixed versus arbitrary intervals
- § Model based on simple scaling properties (power laws) for particular ranges of k values
- § Climatological and hydro-meteorological data
- § Regional model of the at-site means depending on a single at-site characteristic and regional slopes values



9 IMSC

Modeling the at-site means

$$\hat{\mu}_{As}(k) = \begin{cases} \frac{1}{k_0^{c_0-c_1}} \frac{\hat{\mu}_{Fs}(k_1) \cdot H(k_1)}{k_1^{c_1}} \cdot k^{c_0} & (\text{for } k \leq k_0) \\ \frac{\hat{\mu}_{Fs}(k_1) \cdot H(k_1)}{k_1^c} \cdot k^c & (\text{for } k > k_0, \text{ and } c = c_1 \text{ for } k_0 < k \leq k_1, c = c_2 \text{ for } k_1 < k) \end{cases}$$

- § Model based on power laws per range of k
- § integrating the ratio of the arbitrary (A) versus fixed (F) intervals (regional Hershfield factors)

9 IMSC



Modeling the at-site means

- § the at-site means are depending on a single at-site characteristic
- § 2 angular points $k_0=1.028$ and $k_1=5$
- § 3 regional characteristic slopes $c_0=0.2640$, $c_1=0.3544$ and $c_2=0.4688$

9 IMSC





Practical results: Intensity-Duration-Frequency curves and Reduced I.D.F.

$$\hat{I}_{kTs} = \frac{\hat{Q}_{kTs}}{k} = \frac{\hat{\mu}_{ks} \cdot \hat{X}_{kTs}^R}{k} = \hat{\mu}_{Fs}(5) \cdot \hat{R}_{kTs}$$

- § Practical information for specialized customers
- § Useful information for answering to the public and authorities
- § Useful to assess to frequency/return period of particular events
- § Extend the range of k values
- § Regional Reduced I.D.F.

9 IMSC






Conclusions

- § Regional procedures have been adopted for
 - 1 assessment of the regional growth curves
 - 1 Assessment of the scaling properties
 multi-fractal model can describe the evolution of the extreme means
- § Reduced regional I.D.F. curves
- § All the fractiles are obtained by scaling it with the at-site 5-day extreme values means

A regional Intensity-Duration-Frequency curves based on precipitation scaling properties in Belgium

D. Gellens



Regional GEV distribution parameters

$$\hat{\theta} = 7.8590.c + 2.9554.c^2 \quad \text{with} \quad c = \frac{2}{3 + \tau_3} - \frac{\ln 2}{\ln 3}$$

$$\hat{\alpha} = \frac{\lambda_2 \hat{\theta}}{(1 - 2^{-\hat{\theta}}) \Gamma(1 + \hat{\theta})}$$

$$\hat{\xi} = \lambda_1 - \hat{\alpha}(1 - \Gamma(1 + \hat{\theta})) / \hat{\theta} \quad \text{as} \quad \lambda_1 = 1 \quad \hat{\xi} = 1 + \hat{\alpha}(\Gamma(1 + \hat{\theta}) - 1) / \hat{\theta}$$

- § $\lambda_1 = \beta_0 = 1$, the mean; $\lambda_2 = \tau$ (L-moments)
- § β_i are the PWM moments and λ_i are the L-moment linear combination of the β_i
- § $\lambda_2 = 2\beta_1 - \beta_0$
- § $\lambda_3 = 6\beta_2 - 6\beta_1 + \beta_0$

Regional GEV distribution parameters

$$\beta_r = E[X(F(X))^r]$$

The unbiased estimator

$$b_r = \frac{1}{n} \sum_{j=1}^n \frac{(j-1)(j-2)\dots(j-r)}{(n-1)(n-2)\dots(n-r)} X_j$$

where the X_j are the ranked observations

Data extension procedure combined with the regional approach

- § Rather short reference period 1951-1995
- § Few “historical” reference station
- § Possible temporal inhomogeneities due to record interruptions and a.o.b.
- § Method of the fractiles based on the regional GEV distribution to fill the “gaps” and extend the data set to the reference period 1910-1995 (Gellens, J. of Hydrol., 2002)
- § Efficient for mean correlation larger than 0.5

9 IMSC




Modeling the I.D.F. curves and the extreme means

- § Purpose:
 - 1 express the fractiles as a function of k and extend the range of k values
 - 1 combine information from the two networks
- § Modeling the growth curves parameters
- § Modeling the at-site means
- § I.D.F.
- § Regional reduced I.D.F.

9 IMSC




Verification on the location parameter ξ



9 IMSC



Regional growth curves



Regional reduced I.D.F.

$$\hat{I}_{kTs} = \frac{\hat{\mu}_{ks} \cdot \hat{x}_{kT}}{k} = \hat{\mu}_{Fs}(5) \cdot \frac{H(5)}{5^c} \cdot k^c \cdot \frac{\hat{x}_{kT}}{k} = \hat{\mu}_{Fs}(5) \cdot \hat{R}_{kT}$$

- § Separation of the at-site and regional factors
- § Practical objective

9 IMSC

